


September 2012

Oil & Gas Water Use in Texas: Update to the 2011 Mining Water Use Report



Prepared for
Texas Oil & Gas Association, Austin, Texas

Bureau of Economic Geology
Scott W. Tinker, Director
Jackson School of Geosciences
The University of Texas at Austin
Austin, Texas 78713-8924

Oil & Gas Water Use in Texas: Update to the 2011 Mining Water Use Report

**Jean-Philippe Nicot, P.E., P.G., Robert C. Reedy, P.G.,
Ruth A. Costley, and Yun Huang, P.E.**



A handwritten signature in black ink, appearing to read "Jean-Philippe Nicot", written over a faint circular stamp.

**Bureau of Economic Geology
Scott W. Tinker, Director**
Jackson School of Geosciences
The University of Texas at Austin
Austin, Texas 78713-8924

Executive Summary

In Spring 2012, we undertook an update of the hydraulic fracturing sections of the TWDB-sponsored report titled “Current and Projected Water Use in the Texas Mining and Oil and Gas Industry” that we published in June 2011 (Nicot et al., 2011). The 2011 report provided estimated county-level water use in the oil and gas industry in 2008 and projections to 2060. This 2012 update was prompted by two main events: (1) a major shift of the oil and gas industry from gas to oil production, displacing production centers across the state and impacting county-level amounts; (2) rapid development of technological advances, resulting in more common reuse and in the ability to use more brackish water. The timely update was enabled by a faster than anticipated development, translating into abundant statistical data sets from which to derive projections, and by an increased willingness of the industry to participate in providing detailed information about water use in its operations. This document follows the same methodology as the 2011 report but differs from it in two ways. Our current update clearly distinguishes between water use and water consumption. The 2011 report does not include reuse from neighboring hydraulic fracturing jobs, recycling from other industry operations or other treatment plants, and use of brackish water. Our update also presents three scenarios: high, low, and most likely water use and consumption with a focus on water consumption. This update has been reviewed by the TWDB and should supersede oil and gas industry projections from the 2011 report.




Figure ES1. Spatial distribution of hydraulic fracturing water use in 2008 (~36,000 AF) and 2011 (~81,500 AF).

Overall we find that, if the total water use for hydraulic fracturing has increased from 36,000 AF in 2008 to ~81,500 AF in 2011 (Figure ES1), the amount of recycling/reuse and the use of brackish water have also increased (~17,000 AF in 2011, or 21%). Hydraulic fracturing has expanded to the southern and western, drier parts of the state and, by necessity, the industry has had to adapt to those new conditions. Collected information tends to suggest that the industry has

been decreasing its fresh-water consumption despite the increase in water use. Total water use information is relatively easy to access (through the private database vendor IHS), but true consumption is harder to gauge.

The updated hydraulic fracturing projections at the state level do not show a major departure from and are essentially consistent with the previous report but have a more subdued peak and a longer tail (Figure ES2). This is due to the increased likelihood that the industry has hydraulically fractured more formations that can be placed into the tight oil and gas category. The annual peak water use previously estimated at 145,000 AF in the early 2020's is now thought to be a broad peak plateauing at ~125,000 AF/yr during the 2020's. However, fresh water consumption is estimated to stay at the general level of ~70,000 AF/yr and to decrease in future decades. Adding other oil and gas industry water uses, such as waterflooding and drilling, brings projected maximum water use up to ~180,000 AF/yr during the 2020-2030 decade with a much lower consumption which brings the total mining water use to a maximum of ~340,000 AF/yr around the year 2030. These values remain small compared to the state water use (Figure ES3). In 2010, hydraulic fracturing water use represented about 0.5% of the water use in the state. However, the hydraulic fracturing water use is unevenly distributed across the state and may represent locally a higher fraction of the total water use.




Figure ES2. State-level projections to 2060 of hydraulic fracturing water use and fresh-water consumption and comparison to earlier water projections.




Figure ES3. Average state level water use (all categories) in 2001-2010.

Table of Contents

Executive Summary	i
Table of Contents	v
List of Figures	vii
List of Tables	xi
Acknowledgments	xiii
Acronyms	xiii
I. Introduction	1
II. Methodology	3
II-1. Historical and Current Water Use	3
II-1-1 Indicator for Quality Control	3
II-1-2 Hydraulically-fractured Length	5
II-1-3 Beyond the Database	6
II-2. Future Water Use Projections	7
II-3. Notes on Collected Information	8
III. Historical and Current Water Use	11
III-1. Play Description	11
III-1-1 Barnett Shale	11
III-1-2 Eagle Ford Shale	12
III-1-3 TX-Haynesville Shale and East Texas Basin	12
III-1-4 Permian Basin	13
III-1-5 Anadarko Basin	14
III-1-6 East Texas Basin	14
III-1-7 Gulf Coast Texas	15
III-2. Current Water Consumption and Sources	54
III-2-1 Information about Recycling/Reuse and Brackish Water Use	54
III-2-2 2011 HF Water Use and Consumption	54
III-3. Comparison to Earlier Findings	59
III-4. Drilling Water Use	63
IV. Water Use Projections	65
V. Conclusions	93
VI. References	95
Appendix 1: Revision to 2011 Report	97

List of Figures

Figure 1. Comparison of five approaches to computing lateral length (Barnett Shale play).....9

Figure 2. Histograms of lateral lengths according to various approaches: (a) “test”; (b) “survey”; (c) “GIS” (Barnett Shale play); and (d) “test” (Eagle Ford Shale play).....9

Figure 3. Water use intensity in the Barnett Shale play, showing comparison among between top operators in the play.....10

Figure 4. Barnett Shale horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....16

Figure 5. Barnett Shale horizontal water use intensity as a function of (a) depth; (b) operator and depth; and (c) formation thickness.....17

Figure 6. Barnett Shale spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).18

Figure 7. Barnett Shale county-level average lateral spacing.....19

Figure 8. Map view of lateral expression of horizontal wells in the Barnett Shale centered on Tarrant County.....20

Figure 9. Annual well count in Johnson (a) and Tarrant (b) counties.21

Figure 10. Eagle Ford horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....22

Figure 11. Eagle Ford Shale horizontal wells’ water use intensity as a function of (a) depth; and (b) formation thickness.23

Figure 12. Eagle Ford Shale spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).24

Figure 13. Eagle Ford Shale county-level average lateral spacing.....25

Figure 14. TX-Haynesville Shale horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....26

Figure 15. TX-Haynesville Shale horizontal water use intensity as a function of (a) depth; and (b) formation thickness.27

Figure 16. TX-Haynesville Shale spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).28

Figure 17. TX-Haynesville Shale county-level average lateral spacing.....29

Figure 18. Wolfberry verticals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median vertical productive section length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....30

Figure 19. Wolfcamp horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....	31
Figure 20. Canyon Sand horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....	32
Figure 21. Clearfork verticals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median vertical productive section length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....	33
Figure 22. San Andres-Grayburg verticals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median vertical productive section length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....	34
Figure 23. San Andres-Grayburg horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....	35
Figure 24. Permian Basin spatial distribution of water intensity for (a) vertical and (b) horizontal wells.....	36
Figure 25. Permian Basin spatial distribution of (a) vertical well density and (b) density of lateral (cumulative length per area) for horizontal wells.....	37
Figure 26. Permian Basin county-level average lateral spacing.....	38
Figure 27. Granite Wash horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....	39
Figure 28. Cleveland horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....	40
Figure 29. Marmaton horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....	41
Figure 30. Granite Wash horizontal water use intensity as a function of depth.....	42
Figure 31. Cleveland horizontal water use intensity as a function of depth.....	42
Figure 32. Granite Wash spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).....	43

Figure 33. Granite Wash horizontals county-level average lateral spacing.....	44
Figure 34. Cleveland spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).	45
Figure 35. Marmaton spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).	46
Figure 36. Anadarko spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).	47
Figure 37. Map view of wells' lateral expression and vertical well location in the Anadarko Basin.	48
Figure 38. Cotton Valley verticals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median vertical productive section length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....	49
Figure 39. Cotton Valley horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....	50
Figure 40. Cotton Valley horizontal water use intensity as a function of depth.....	51
Figure 41. Cotton Valley spatial distribution of density of lateral (cumulative length per area).....	51
Figure 42. Cotton Valley spatial distribution of density of vertical wells (years 2005-2011).....	52
Figure 43. Olmos horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.....	53
Figure 44. Location of waste water treatment facilities that provide or have provided water to the industry for HF as of July 2012.	58
Figure 45. Spatial distribution of HF water use in 2008 and 2011.	60
Figure 46. Bar plot comparison of 2011 actual water use to projections from 2009.....	61
Figure 47. County-level cross-plot comparison of 2011 actual water use to projections from 2008. Values on x- and y- axis represent counties whose actual (y-axis) / projected (x-axis) water use is 0. A total of 168 counties are represented.	62
Figure 48. State-level projections to 2060 of HF water use and fresh-water consumption and comparison to earlier water projections.	81
Figure 49. State-level projections to 2060 of oil and gas industry water use and fresh-water consumption.	81
Figure 50. Counties with non-zero projected water use. Same coverage as in the 2011 report (thick blue lines) with the addition of Polk, Tyler, Jasper, and Newton counties in East Texas (red circle).	82
Figure 51. Spatial location of the oil and gas windows in the (a) Barnett Shale and (b) Eagle Ford Shale.	82
Figure 52. Barnett Shale water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.....	83

Figure 53. Eagle Ford Shale water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.....	84
Figure 54. Pearsall Shale water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.....	85
Figure 55. Haynesville and Bossier Shales water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.....	86
Figure 56. Haynesville-West Shale water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.....	87
Figure 57. East Texas (not including Haynesville and Bossier Shales) water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.	88
Figure 58. Gulf Coast (not including shales) water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.....	89
Figure 59. Anadarko Basin water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.....	90
Figure 60. Permian Basin water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.....	91
Figure 61. Permian Basin water use and consumption projections under the three scenarios: (a) Barnett and Woodford Shales; (b) Wolfcamp Shale and Wolfberry play; and (c) other Permian Basin formations.	92
Figure 62. Summary of projected water use by mining industry in Texas (2012-2060).	94
Figure 63. Average state level water use (all categories) in 2001-2010.....	94

List of Tables

Table 1. Representivity of collected information	10
Table 2. Percentage of wells in each play or region that yielded a complete and consistent data set (water, proppant, length) from year 2011.	15
Table 3. Barnett Shale county-level average lateral spacing for top producing counties.	19
Table 4. Eagle Ford Shale county-level average lateral spacing for top producing counties.	25
Table 5. TX-Haynesville Shale county-level average lateral spacing for top producing counties.	29
Table 6. Granite Wash county-level average lateral spacing for top producing counties	44
Table 7. Estimated percentages of recycling/ reused and brackish water use in main HF areas in 2011.	56
Table 8. Estimated groundwater / surface water split (does not include recycling / reuse)	56
Table 9. HF water use in 2008 and 2011 compared to the 2011 projected water use from 2008.	56
Table 10. County-level estimate of 2011 HF water use and water consumption (kAF).	57
Table 11. Drilling water use information.	63
Table 12. Recent trends in well completion and water use in hydraulic-fractured plays.	69
Table 13. Coefficients (%) to compute water consumption to be applied to total water use.	69
Table 14. Estimated flow back/produced water volume relative to HF injected volume.	70
Table 15. County-level estimate of 2012-2060 projections for HF water use and water consumption (AF).	71
Table 16. County-level estimate of 2012-2060 projections for oil and gas water use and water consumption (AF).	76
Table 17. Update to Table 52 of 2011 report (now obsolete and superseded by this report)	97

Acknowledgments

The authors would like to thank engineers and staff from many oil and gas operating companies for giving their time in answering our requests and sharing their knowledge of the hydraulic fracturing process. Special thanks to C. J. Tredway for managing the project, to Dr. Dan Hardin (TWDB) and his staff for reviewing the report, and to BEG internal reviewers, particularly Dr. Bridget Scanlon. Thanks too to L’Oreal Stepney and her staff at TCEQ for providing useful information. We are also grateful to IHS for providing free access to their Enerdeq database, which was used extensively in the course of this work. The report also benefited from a thorough editing by Amanda R. Masterson of the Bureau of Economic Geology at The University of Texas at Austin.

Acronyms

AF	Acre-foot
BEG	Bureau of Economic Geology
EOR	Enhanced Oil Recovery
Fm.	Formation
GW	Groundwater
HF	Hydraulic fracturing
kAF	Thousand acre-feet
Mgal	Million gallons
PSD	Powell Shale Digest
RRC	Railroad Commission (of Texas)
SW	Surface water
TCEQ	Texas Commission on Environmental Quality
TDS	Total dissolved solids
TWDB	Texas Water Development Board
TXOGA	Texas Oil & Gas Association

I. Introduction

This work is an update of the “Current and Projected Water Use in the Texas Mining and Oil and Gas Industry” (Nicot et al., 2011) report released in 2011 by the Texas Water Development Board (TWDB) and prepared by the Bureau of Economic Geology (BEG). The 2011 report documents future and projected water use in all segments of the mining industry: oil and gas, aggregates, coal, and other industrial and metallic substances. In particular, it looked at three main water categories in the upstream segment of the oil and gas industry: drilling, waterflooding and enhanced oil recovery (EOR), and hydraulic fracturing (HF).

How is this report different from the 2011 Report?

This report focuses on HF water use and associated drilling; the information in the 2011 report relating to waterflooding and EOR water use as well as drilling not associated with hydraulically-fractured wells did not require updating. This update also benefited from more participation from the industry, especially for information not typically available or easily extractable from state records. We also have a longer record for many plays, indicating trends and allowing for better future projections. In addition, we presented three scenarios for water use and water consumption for each play (high, medium, low) as was done in Bené et al. (2007) but not in the 2011 report. Furthermore we made the distinction between water use and water consumption more explicit. Water use is the amount of water used in an operation regardless of the water source provided; water is either fresh or brackish. Fresh water is defined as any water with a total dissolved solids (TDS) content of <1,000 mg/L; the upper limit for brackish water is 35,000 mg/L, but often in this document the limit will be <10,000 mg/L. Water consumption is fresh water use excluding recycling and reuse. Reuse is understood as the water originating from previous HF operations whereas recycling is more general and could include, for example, produced water from conventional wells or waste water obtained from other industries or municipalities.

Scope of work

As in the 2011 report, this update’s scope of work includes two main tasks: (1) documenting current (year 2011) and past water use from HF; and (2) estimating projected water use. Both tasks are completed at the county level for the entire state of Texas. Task 1 consists of gathering water use data and establishing statistics needed for the projection phase in the spirit of what was done in the 2011 report but with a more detailed processing of the data. Task 2 is to produce a projection of county- level water use to 2060 using previously derived statistics and input from the industry.

This current document is organized in the following way. We first describe the methodology and its caveats as well as the challenges to making projections. We then examine the 2011 water use and compare our new findings to the 2011 projections made in 2008 as a way to validate our approach. We then present projections to 2060 according to three scenarios: high estimates, most likely estimates, and low estimates.

II. Methodology

II-1. Historical and Current Water Use

We followed a methodology similar to that used in the 2011 report, making use of the IHS Enerdeq database (<http://www.ih.com/products/oil-gas-information/data-access/enerdeq/browser.aspx>). The IHS data were cross-checked with information from individual companies (number of oil/gas wells, of vertical/horizontal wells, amount of proppant) through discussion with company experts. In addition to production data, the Enerdeq database contains completion information submitted by operators to the Railroad Commission (RRC) of Texas through the W-2 and G-1 forms for oil and gas, respectively. In the best cases, and as noted by statistics provided in forthcoming sections of this report, the database contains all information of interest to us: API number, location of the well, well geometry, amount of water used, and amount of proppant used. Because, across plays, the completeness of the data is variable and because typographical errors are not infrequent, we developed several indicators for quality control: water intensity (amount of water used per unit length of lateral or useful vertical section) and proppant loading (amount of proppant per unit water volume). When either water intensity or proppant loading for a given well is out of range, the well is flagged and obvious errors corrected (for example, reporting water use in gal but displaying bbl as the unit instead of gal). Details on the approach follow.

The three primary data types used to estimate HF water volumes include reported values of fluid and proppant used to fracture each well and the total well length over which fracturing procedures were performed. Data were extracted separately from the IHS database for individual producing formations having a significant number ($> \sim 100$ to 200) of wells located in Texas that were completed between January 1, 2005 and December 31, 2011 that upon preliminary accounting had been fractured using $> 100,000$ gal of fluids. These include the Barnett, Eagle Ford, Haynesville, Cotton Valley, and Olmos formations, and several formations in the Anadarko Basin (Granite Wash, Cleveland, Marmaton) and the Permian Basin (Wolfcamp, Spraberry, Canyon, Clear Fork, San Andres, and Grayburg). For this analysis, the Wolfcamp and Spraberry were combined and the San Andres and Grayburg were combined.

As we did in the 2011 report we relied on the IHS database to recognize the currently active plays by downloading basic information on all wells drilled in Texas since 2010 (included early 2012 but with many gaps in the reporting). Our interest was not in computing water use but in determining those plays with enough activity to warrant a more detailed study. Many additional wells were fractured in other plays and did count toward the total water use in 2011, but they were not part of the detailed analyses of those plays cited earlier. Those minor plays are, however, accounted for in the general Gulf Coast and Permian Basin count.

II-1-1 Indicator for Quality Control

For producing formations having a sufficient number of wells completed during this period, the data were analyzed by annual intervals. Wells having actual or estimated total HF water use of $< 100,000$ gal (i.e., small-scale traditional fracturing performed primarily on vertical/directional wells) were omitted from calculations as they account for comparatively insignificant water volumes compared to the fracturing currently being practiced in many plays. This minimum

volume distinction was applied to vertical/directional wells only, and all horizontal wells were included in the estimates.

Critical evaluation and editing of the raw data was required. The purpose of the editing process was, through a step-wise logical procedure, to exclude wells that used or (in the absence of accurate data) were likely to have used <100,000 gal of HF fluids while retaining and accounting for wells that used or (again, in the absence of accurate data) were likely to have used $\geq 100,000$ gal of HF fluids. For many wells, one or more of the reported data values is absent, incomplete, or inaccurate, due either to clerical errors or to partial reporting (omission errors). Clerical errors include the incorrect assignment of units (gal vs. bbl, lb vs. ton, etc.) and/or typographical errors. Omission errors primarily include the non-reporting or under-reporting of fluid volumes (proppant amounts seem to be accurately reported much more consistently than fluid volumes).

The data were screened for errors by examining ratios between the different values, including the total reported volume of fluids used per linear foot of the total fractured well depth interval (water use intensity, gal/ft), the total mass of proppant per total volume of HF fluids (proppant loading, lb/gal), and the total mass of proppant per linear foot of the total fractured well depth interval (proppant intensity, lb/ft). These ratios were examined for outliers and inaccuracies by sorting hierarchically through the data based on the various ratios. Edits were performed on the raw data where rectifiable errors could be identified, the most prevalent consisting of modifying units where such changes resulted in ratios consistent with other similar wells. In some cases, sufficient details were reported in the data comments to correct inaccurate data values, although this type of edit was extremely limited.

In general, proppant loading (lb/gal) was used as the primary data screening ratio because of the generally consistent reporting of total proppant amounts. HF fluid volumes resulting in proppant loading values (average of all stages) > 5 lb/gal were deemed as under-reported. Barring a unit's error, these values generally reflect reported fluid volumes that include only acid treatments and in some cases raw gel product volumes and do not also include the volumes of water used. For vertical/directional wells having reported proppant amounts and with absent or under-reported HF volumes, wells with <100,000 lb of proppant were excluded from the estimates based on an assumed 1.0 lb/gal loading ratio.

A finer level of resolution in the water use data could be achieved by binning the hydraulic fracturing stages into slickwater, gel, and cross-linked gel systems with the latter two having a smaller water use intensity. Unfortunately the database does not allow for an accurate count in each category. The information, however, was used in a qualitative way, checking its consistency with common practices in a play.

Following the data screening and editing procedures, the data were classified into two main groups: 1) wells judged to have accurately reported fluid volumes and 2) wells judged to have inaccurately reported fluid volumes. The average (annual) water use intensity (gal/ft) values of the Group 1 wells were multiplied by the (annual) sum total fractured length (ft) of the Group 2 wells to produce annual estimates of the total water use of the Group 2 wells. The average intensity values represent truncated averages based on 90% of the data that were calculated by eliminating values less than the 5th percentile or greater than the 95th percentile of the Group 1 population to reduce the impacts of extreme values. The Group 2 annual total estimates were then added to the Group 1 annual total values to produce estimates of actual annual total water

use. Values are reported for the major producing formations listed above by year and by county. County locations were assigned based on the wellhead coordinates.

A separate estimate using the same procedures was calculated for the HF water used during 2011 for all wells meeting the minimum 100,000 gal criteria but that were not completed in one of the producing formations listed above and for which insufficient data exist for temporal trend analysis.

II-1-2 Hydraulically-fractured Length

HF lengths for individual wells were determined using five approaches, each relying on different information in the database. All five approaches were applied to varying degrees to determine horizontal well HF lengths while only the first two were applied to vertical/directional wells. The first approach used the difference between the minimum and maximum reported test treatment depths and is referred to as the “test” length. This was the primary length used in an estimated minimum of 95% of all wells. The second approach used the difference between the minimum and maximum perforation depths, which was identical in most cases to that of the test length and is referred to as the “perf” length. The “perf” length was used in place of the test length in a few cases that resulted in more realistic use intensity values. The test and “perf” lengths are considered to be the most accurate length information available for most wells.

A third approach utilized the survey information and is referred to as the “survey” length. In this approach, the angle relative to the horizontal plane between successive well survey points was calculated. The horizontal length of the well was determined as the difference between the minimum depth at which that angle became less than 2.5 degrees and the maximum well depth. This approach also provided the average depth of the horizontal well section and additionally the beginning and ending X-Y coordinate locations of the horizontal well section used to map well density in GIS for the various plays. If no information was available to calculate a test or perf length, the survey length was considered to be the next-best available length information. In most cases where all three were available, the survey length is in good agreement with both the test and perf lengths. This value was used only in a few cases where neither a test nor a perf length was available.

A fourth length value was calculated as the difference between the reported driller’s well depth and the bottom hole true depth, referred to as the “true value” or “TV” length and a fifth length value was calculated as the simple horizontal linear distance between the X-Y coordinates of the well surface and bottom hole coordinates (“GIS” length). Both of these values are considered to be only general estimates of the horizontal section length and were used in a very limited number of instances where more accurate information was not available. For a very few instances (<<1%) no length values were available for a given well. In these cases, the annual (truncated) average well length for that producing formation was assigned.

The fourth and fifth approaches, simpler to use, were adopted in the 2011 report. The HF water intensity for horizontal wells is computed slightly differently from the approach in the 2011 report. Instead of using the distance between the wellhead of the toe of the lateral, we used a shorter distance defined by the operator-defined “test length” more representative of the true length of the lateral. The test length is consistent with the “test” length but consistently smaller by 10 to 25%. The lateral length value matters as it used to compute water intensity, itself used to make projections. There is relatively little difference between the different approaches (Figure 1)

but the “test” approach used in this document is systematically smaller than the “GIS” approach used in the 2011 document, that is, water intensity values reported in this document are systematically greater than those in the 2011 report. The median value of water intensity using the “test” and “survey” approaches are 26% and 23% larger than the “GIS” median value (Figure 2) in the Barnett Shale play. The “test” water intensity median in the Eagle Ford play is 16% larger than the “GIS” median value (Figure 2d).

II-1-3 Beyond the Database

In the 2011 report we made the explicit distinction between shale plays and tight gas plays. Although, as explained in the 2011 report, there are real differences between them, from an operational standpoint the difference is blurred (for example, wells tapping Wolfcamp shale oil and Spraberry tight oil) and, in this update, we did not try systematically to assign one of either category to some plays.

For each of the plays with sufficient data we extracted yearly information, presented in the Results Section, about:

- Total number of wells
- Total water use, including estimation of data gaps
- Average/median length of laterals
- Water use in Mgal/ft
- Water intensity in gal/ft
- Proppant loading in lb/gal

The IHS database provides only water use, that is, the amount of water used during a given HF job regardless of the water source(s). In actuality, water can come from several sources. It can be “new” water or it can also be recycled or reused water. “New” water can be surface water or groundwater or it can be from an alternative source such as municipal water or treated waste water. Water also be fresh (<1,000 mg/L) and its use can directly compete with other more conventional users (municipal use, irrigation use). It can be brackish or even more saline than sea water (that is, >~35,000 mg/L). Water consumption is simply defined as the water use which is not from recycled or reused water and from which brackish and saline water use is taken out. Note, however, that this simple definition does not capture a more complex reality. Use of brackish water in areas with limited fresh water supplies could compete with conventional users. This document does not try to sort out such issues; we simply define water consumption as water use minus recycled/reused water volumes and minus brackish or saline water volumes.

Access to detailed information about water sources on the provider side is difficult. Large water suppliers do not necessarily track the ultimate usage of their water. Groundwater conservation districts (GCD’s) do not always collect information about withdrawal amounts and eventual use of the water. A request to the Texas Commission on Environmental Quality (TCEQ) on reuse of treatment water yielded a helpful list of facilities but not the amount of water transferred, and further this does not account for direct reuse at a site. The demand side, that is, operators, is very fragmented.

We collected information not present in the IHS database but of interest to TWDB and the general public about: (1) nature of the water source (river, lake, city water, groundwater, stock pond/gravel pit / quarry, wholesaler, treated industrial waste water) and its status (private, public). The ultimate goal is to determine the groundwater and surface water (GW/SW) split. Optimally,

this issue would be resolved at the county level but it may not be possible; (2) amount of water injected from reuse of flow back water, recycled water can include water from commercial and municipal waste water treatment facilities; (3) TDS of the new water [fresh (<1000 mg/L), slightly brackish (1000-3000 mg/L), brackish (3000-10,000 mg/L or 10,000-35,000 mg/L), saline (>35,000 mg/L)].

In this document, we applied to all counties within a play / region the same brackish water use, recycling/reuse fraction, and GW/SW split. Undoubtedly, this is an approximation but the amount of information available does not allow accurate assessments at the county level.

II-2. Future Water Use Projections

The 2011 report followed a mixed approach to estimate projected water use, the so-called resource-based and production-based approaches. Although both approaches are somehow interdependent, we believe that the resource-based approach gives the best results and is used in this document. As described in more details in the 2011 report, it consists of four steps:

- (1) Gather historical data in terms of average well water use and average well spacing. It is important to establish these elements through time to see trends rather than just focusing on the past few months.
- (2) Estimate ultimate well density across the play; it is a function of several factors, such as geological prospectivity (for example, within play core or not, shale thickness) and cultural features (urban/rural). In this step, ultimate boundaries of the play are identified.
- (3) Compute approximate total number of wells needed.
- (4) Distribute through time and space, constrained by the assumed number of drilling rigs available (see earlier comment).


After obtaining water use, correction factors to account for recycling/reuse and use of non-fresh water are applied. We asked industry operators for projected recycling/reuse, brackish water use, and groundwater / surface water split in 2020. Given the rapid pace of change in the industry, the values obtained are somewhat speculative. Although not a guarantee for accuracy, those values are, however, consistent with what industry observers report and consistent with our own knowledge of treatment techniques and state of surface water and groundwater withdrawals across the state. The basic reporting unit for the water use projections is the county. Projections for recycling / reuse, brackish water use beyond 2020 to 2060, were made accounting for the typical current volume of flow back (limiting reuse) and for brackish water resources / lack of fresh water in the area of interest.

As discussed in the 2011 report, despite our best efforts, it is likely that the projected water use amounts will be more accurate at the play than at the county level. As done in the 2011 report, we did not assume any repeat HF, as discussions with industry experts and recent publications (Sinha and Ramakrishnan, 2011) suggest that little repeat HF will take place.

The 2011 report provides only one annual estimate. However, in an earlier report on the Barnett Shale only (Nicot and Potter, 2007; Bené et al, 2007), BEG made use of high, medium, and low water use scenarios. The different scenarios were based on various level of prospectivity and anticipated gas price. This update also makes use of three scenarios, high, most likely, and low water use, but in addition to prospectivity and gas price, they take into account level of recycling/reuse and use of brackish and saline water.


II-3. Notes on Collected Information

We obtained information on all the major plays, some with better coverage, by contacting operators. Fraction of HF wells drilled by contacted operators in the 2010-2012 period is documented by play and provides an estimate of the uncertainty. The coverage (Table 1) was calculated by adding the number of wells completed in the 2010-early 2012 period by contacted operators and normalizing that sum by the total number of wells completed during the same period. We collected information about recycling/reuse, use of brackish water, surface water/groundwater split. Coverage varies from 40% (Barnett Shale) to 10.5% (Permian Far West). Consistency in information from operators in a given play suggests that even low percentages are representative of the industry as a whole in that play despite some variability among operators (Figure 3). The figure shows a slight overall increase in water use intensity with increasing depth but it also shows that operators can have different approaches.



Histogram2.0 lateralComputation Barnett.xlsx

Figure 1. Comparison of five approaches to computing lateral length (Barnett Shale play).



Histogram2.0 lateralComputation Barnett.xlsx

Histogram2.0 lateralComputation EagleFord.xls

Figure 2. Histograms of lateral lengths according to various approaches: (a) “test”; (b) “survey”; (c) “GIS” (Barnett Shale play); and (d) “test” (Eagle Ford Shale play).

Table 1. Representivity of collected information

Play/Region	Consumption information (%)
Permian Far West	10.5%
Permian Midland	23%
Anadarko Basin	11%
Barnett Shale	40%
Eagle Ford Shale	31.2%
East Texas Basin	14.5%
All Plays	27.2%




Figure 3. Water use intensity in the Barnett Shale play, showing comparison among between top operators in the play.

III. Historical and Current Water Use

After a short description of the major HF plays in Texas (Section III-1), we present water use and consumption numbers (Section III-2) that we compare to findings of the 2011 report (Section III-3). We also briefly address drilling water use (Section III-4).

III-1. Play Description

In this section we describe relevant features of each play which will then be used in the Projections Section (Section IV). Note that water use intensity and proppant loading values represent an average of the sometimes time-varying mix of slickwater / gel systems applied to the play at a given time. For example, a decrease in water use intensity may mean a better water efficiency in a technique or a move to a more water-efficient technique.

III-1-1 Barnett Shale

The Barnett Shale is the first in Texas and around the world to submit to intense slick-water HF since the mid-1990's, first using vertical wells. After a transition period, Barnett Shale operators use currently horizontal wells almost exclusively. After a strong growth in the mid-2000's (>2000 wells completed per year), the play has seen a relative decrease in the total number of wells completed in a year (Figure 4a) because of the reduced demand following the economic slump and the decreasing price of gas. Although drilling activity has abated at the edges of the play core, it is very vigorous in the core itself (Denton, Johnson, Tarrant, and Wise counties) and has considerably picked up in the so-called combo play in the northern confines of the play in Cooke and Montague counties. A weekly newsletter, the Powell Shale Digest (PSD; May 29, 2012) noted a sharp increase in oil production since mid-2010. Substantial amounts of oil and condensate have made those counties attractive to operators. Overall the total amount of water used is relatively steady at 25 kAF/yr (Figure 4b). The Barnett play is the Texas play with the highest degree of reporting water use at >90% (Table 2). Note that the bottom four plots of composite Figure 4 (as well as on similar figures in this document) show the fraction of wells used to compute the parameter on the secondary axis. High well reporting, allied with the large number of wells, gives us confidence that the water use values are particularly accurate in this play. The length of the laterals has been slowly increasing in the past few years (~3,500 ft in 2011) with a concomitant water use increase (Figure 4c and d). However water intensity (water amount per unit length) has stayed steady at ~1,200 gal/ft (Figure 4e). Note that the water intensity as reported in this document is higher than that reported in the 2011 report because of a slight change in computing it (see Section II-1-2). In contrast to water intensity, proppant loading has been increasing slightly over time to ~0.8 lb/gal in 2011 (Figure 4f).

In order to better understand water intensity and in an effort to modulate it across a play, we plotted water intensity against depth and thickness (Figure 5a and c). The trend seems upwards with increasing depth and thickness but is very noisy and tenuous at best. Water intensity appears to be rather dependent on the well operator (Figure 5b) and, thus, somehow difficult to vary across a play. Nevertheless, spatial distribution of water intensity shows a higher intensity in Denton County and in the eastern half of Wise County, areas in which the Barnett is the deepest as well as in Montague County in the oil window (Figure 6a).

In agreement with our methodology, it is also useful to understand the cumulative length of laterals in a given area or within a county. A key input to the projected water use is to assume

that the entire county will be hypothetically drilled up by parallel laterals extending from one side of the county to the other side and at regularly spaced intervals (at, for example, a 1,000- ft interval [see Nicot et al., 2011 for details]). Figure 6b displays such density of well laterals, which is fairly high in Johnson County and the southern half of Tarrant County. The average lateral spacing, which is simply the inverse of the lateral density, is shown in Figure 7 and detailed in Table 3 (it is calculated in those sections of the county with an actual shale footprint). The county with the highest relative cumulative length of laterals (Johnson County) yields an average spacing between assumed parallel laterals of ~1,700 ft. This is still removed from the operational distance between laterals of 1,000 ft or even 500 ft, suggesting that this county, despite its past activity will still see further significant activity as illustrated by the coverage gaps in Figure 8. The decrease in well completion activity in Johnson County as seen in Figure 9a is more related to price gas than to a true depletion of the resource in the county.

III-1-2 Eagle Ford Shale

The Eagle Ford Shale play has seen tremendous development in the past 2 years. Initially started as a new Barnett Shale, it quickly turned into a different type of play when the extent of the oil window became clear. In addition to the fast increase in wells completed (~1,400 in 2011) (Figure 10a) and the subsequent increase in water use at ~24 kAF in 2011 (Figure 10b), the Eagle Ford Shale has the unique feature among all the plays examined in this document to experience a sharp decrease in water intensity (Figure 10e) decreasing almost in half in 4 years to ~850 gal/ft in 2011. This is seemingly due to operational changes moving from high-volume slick water HF operations to gel fracs that can carry as much proppant with much less water. The use of cross-link gels for oil production requires a higher proppant loading (Fan et al., 2011). This decrease in water intensity combined with an increase in average lateral length (~5,000 ft, Figure 10c) still translates into a decrease in water use per well to ~5 million gallons/well (Figure 10d). Not surprisingly, the proppant loading has considerably increased to 1 lb/gal in 2011 (Figure 10f). The question we will not try to answer despite its relevance to water use projection is how transferable to other plays is this switch to gel fracs and whether it could happen elsewhere on a large scale. The percentage of wells with consistent data sets is only ~47% (Table 2), making the Eagle Ford data set more uncertain than that of the Barnett Shale.

The cross-plots of water intensity vs. depth and thickness are inconclusive and even misleading (Figure 11a and b). They show no real trend except perhaps a decrease in water intensity with depth. However, Figure 12a clearly shows a higher water intensity in the down dip sections of the play, suggesting an intensity as high as 1400 gal/ft in the gas-rich area and 800 gal/ft in the oil-rich area. Densities of lateral (Figure 12b) and average lateral spacing (Figure 13, Table 4) suggest that the Eagle Ford Shale play has two cores: next to the Mexican border in Dimmit, LaSalle, and Zavala Counties and south of San Antonio in Karnes and De Witt Counties. The low average lateral spacing (>10,000 ft) suggests that many more wells will be drilled and completed there in the future.

III-1-3 TX-Haynesville Shale and East Texas Basin

This document deals only with the Texas section of the Haynesville Shale. In East Texas the Haynesville is a deep gas play, despite a report that one company has located a liquid-rich area in the Haynesville in Panola County with 350 horizontal drill sites (PSD, May 29, 2012). These are expensive wells, but they are located in an area with multiple stacked formations amenable to

HF. The Texas section of the play has seen a quick increase in the number of wells drilled (~250 in 2011, Figure 14a) and a subsequent increase in water use (~1.6 kAF, Figure 14b). This play, with the Cotton Valley Fm., also in East Texas, has the smallest fraction of wells with usable data (32% in 2011, Table 2). Lateral length (~5,00 ft), well water use (~8 million gal/well), and water intensity (~1,400 gal/ft in 2011) have all increased in the past 3 years (Figure 14c, d, and e) whereas proppant loading has stayed stable at 0.8 lb/gal (Figure 14f). Water intensity as a function of depth and thickness does not show any reliable pattern (Figure 15). Water intensity (Figure 16b) and density of lateral (Figure 16c) are spatially correlated. The highest correlations are in Harrison County and where Shelby and San Augustine counties meet (Harrison, Shelby, San Augustine, and Panola counties are all in the TX-Haynesville core area). County-level average lateral spacing (Figure 17 and Table 5) with a minimum value at ~24,000 ft suggests that many more wells will be completed in this play.

III-1-4 Permian Basin

The Permian Basin, comprising the Midland Basin to the East and the Delaware Basin to the West, with the Central Platform in between, has a long history of mostly oil production. It has also received much attention recently because of hydraulically fractured vertical wells in the so-called Wolfberry play (Wolfcamp and Spraberry, Figure 18). More recently, attention has shifted to horizontal wells in the Wolfcamp Shales (Figure 19), one of the source rocks of the many oil accumulations in the Permian Basin. Several other plays are also being hydraulically fractured in the basin such as the Canyon Formation (Figure 20), the Clear Fork Formation (Figure 21), and the San Andres (Figure 22 and Figure 23) among others.

The Wolfberry was the first play in the Permian Basin to benefit from the technological progress made in the Barnett Shale play. The wells are vertical and have grown from <500 wells/yr to >1,500 wells in 2011 (Figure 18a). The annual amount of water use had also increased to almost 8 kAF in 2011 (Figure 18b). Approximately 80% of the wells have consistently good data. As the length of the productive vertical section has increased from 1,500 ft to >2,500 ft in the past few years (Figure 18c), so has the average water use per well which is >1 million gal/well in 2011, relatively small volume compared to that of horizontal wells in shale plays. As productive sections become longer, the water intensity increased slightly to ~400 gal/ft (Figure 18e), but proppant loading remained constant at ~0.9 lb/gal (Figure 18f). Water intensity seems to be higher in the Wolfberry of the Delaware Basin (Figure 24a), but that basin contains very few wells (Figure 25a), (and they might even be misnamed). The well density is the highest in Glasscock and Reagan Counties.

Slick water horizontal wells have been jumped in 2011 from a low level of <50 wells/yr to 160 wells (Figure 19a), with a concomitant increase in total water use (~1.5 kAF in 2011, Figure 19b). Lateral length (~5,000 ft in 2011), well water use (~5 million gal/well in 2011), and water intensity (800 gal/ft in 2011) all increased too (Figure 19c, d, and e), but average proppant loading stayed steady at ~1 lb/gal (Figure 19f). Water intensity is higher in the center of the Midland Basin (Figure 24b), and the density of lateral is the highest in Ward County (Figure 25b) but the average lateral spacing is still very high at ~23,000 ft (Figure 26), which suggests that many wells remain to be drilled and completed.

Other, less publicized plays also received increased interest, as shown by water intensity rising or remaining steady (Figure 20e, Figure 21e, Figure 22e, and Figure 23e). Other plays, not targeted for the same scrutiny, have also seen a development of HF. They were included in a

miscellaneous file that included all fractured wells not included in a targeted play. Overall the Permian Basin has a high fraction (~85%) of wells with a consistent data set (Table 2), thus giving us confidence that the water use values are relatively accurate (especially for those formations hosting a large number of wells).

III-1-5 Anadarko Basin

The Anadarko Basin contains several formations of interest, in particular the Granite Wash (Figure 27) but also the Cleveland and Marmaton formations (Figure 28 and Figure 29). Similarly to the development of the horizontal wells in the Wolfcamp in an area where HF was done on mostly vertical wells, the Anadarko Basin is seeing a shift toward horizontal wells. The Granite Wash has seen an increase from a few horizontal wells in 2006 to >300 in 2011 (Figure 27a) with a parallel increase in water use to <4 kAF in 2011 (Figure 27b). In the same time the length of the lateral has grown to ~4,500 ft (in 2011) (Figure 27c) and the average well water use to >5 million gallons (Figure 27d). Water intensity has reached a value of ~1,200 gal/ft (Figure 27e), but the proppant loading has remained steady at ~0.6 lb/gal (Figure 27f). The Cleveland and Marmaton horizontal wells display a similar evolution but for a smaller number of wells (~150 and ~40, respectively) and smaller water intensity at ~300 gal/ft (Figure 28e and Figure 29e). The fraction of wells with directly usable information was calculated at ~70% (Table 2). Water intensity as a function of depth failed to show a clear trend (Figure 30 and Figure 31).

Spatial distribution of Granite Wash water intensity (Figure 32a) and density of lateral (Figure 32b) confirms that Wheeler County is the most attractive county. At the county level, Wheeler County shows the smallest lateral spacing and plenty of room for additional wells (Figure 33 and Table 6). HF activities in the Cleveland and Marmaton Formations are focused on Hemphill, Lipscomb, and Ochiltree Counties (Figure 34 and Figure 35). Combining information from the three plays illustrates that the county with the smallest average lateral spacing (Lipscomb County) still allows for significant development at ~11,000 ft (Figure 36), as illustrated in Figure 37.

III-1-6 East Texas Basin

The East Texas Basin contains many formations susceptible to being hydraulically fractured. This section focuses on the Cotton Valley Fm., but, as was done for the Permian Basin and the Gulf Coast Basin, all water use data from wells in formations that were not part of the plays targeted for detailed study were still added to the total water use.

The Cotton Valley Fm. has been producing for decades and has been subjected to HF for almost as long. However, as observed in the rest of the state, there is a general shift from vertical to horizontal wells. Annual completions of vertical wells have been decreasing from ~1500 wells per year in 2007 to ~300 in 2011 (Figure 38a), whereas horizontal wells have been increasing from almost none in 2005 to ~100 in 2011 (Figure 39a). Total water use has followed the same path from ~1.5 kAF/yr to ~0 and from ~0 to 0.6 kAF/yr, respectively (Figure 38b and Figure 39b). In 5 years, the length of lateral has increased from ~1,000 ft to ~4,000 ft in 2011 (Figure 39c) with the associated water use increase to 4 million gallons per well in 2011 (Figure 39d). In the same period, water intensity has stayed steady at ~1,000 gal/ft (Figure 39e) and proppant loading has remained at ~0.8 lb/gal (Figure 39f). The overall representivity of the usable data set is at a steady ~70% for the horizontal wells but decreasing to only 25% for the vertical wells. A water intensity vs. depth cross-plot (Figure 40) displays no obvious trends but maps of well

density (Figure 41 and Figure 42) show that horizontal wells are being completed in the same areas as where the vertical wells were drilled and that there is a good overlap of the high density values.

III-1-7 Gulf Coast Texas


Similarly to the Permian Basin and the East Texas Basin, the Gulf Coast Basin, which includes many counties from the Mexican border to the Louisiana state line, contains several formations amenable to being hydraulically fractured. Each of these formations is not described here (for example, the Austin Chalk), but their water use is included in the total reported below. In this section, we document the Olmos Sands, where HF is taking place through horizontal wells. The annual number of completion is still low at 70 completions a year (Figure 43a) but growing and the total water use displays the same growth (~0.5 kAF in 2011, Figure 43b). Average lateral length has reached ~4,000 ft in 2011 (Figure 43c), and the average water use per well has increased to 4 million gal/well (Figure 43d). Although irregular through the years, water intensity has reached a value of ~1,000 gal/ft (Figure 43e) consistent with what has been observed elsewhere.

Table 2. Percentage of wells in each play or region that yielded a complete and consistent data set (water, proppant, length) from year 2011.

Play / Region	Percent
Barnett	92.7%
Eagle Ford	46.9%
Haynesville	31.8%
Cotton Valley	31.4%
Anadarko	69.4%
Permian Basin	84.9%

ResultsSummary_year2011.xlsx


Barnett Shale:




Note: red squares represent average ; blue diamonds represent median; only partial data for 2012

Figure 4. Barnett Shale horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.


Barnett Shale:



(a)





(b)



(c)

Figure 5. Barnett Shale horizontal water use intensity as a function of (a) depth; (b) operator and depth; and (c) formation thickness.

Barnett Shale:



Note: $25 \text{ km}^2 = 154 \times 40$ acres, that is, $154 \text{ wells}/25 \text{ km}^2 = 1 \text{ well}/40$ acres

Figure 6. Barnett Shale spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).

Barnett Shale:




Figure 7. Barnett Shale county-level average lateral spacing.

Table 3. Barnett Shale county-level average lateral spacing for top producing counties.

County Name	Sum lateral length / county area (km/km ²)	Average Lateral Spacing (1000 ft)
Johnson	1.94	1.69
Tarrant	1.66	1.98
Hood	0.75	4.35
Parker	0.53	6.20
Wise	0.48	6.77
Denton	0.47	6.99
Somervell	0.34	9.76
Others		>10×10 ³ ft

Note: Average spacing = 1/ (lateral length density);
 Counties are sorted by decreasing lateral length density






Figure 8. Map view of lateral expression of horizontal wells in the Barnett Shale centered on Tarrant County.




(a)



(b)

Figure 9. Annual well count in Johnson (a) and Tarrant (b) counties.


Eagle Ford Shale:




Note: red squares represent average ; blue diamonds represent median; only partial data for 2012

Figure 10. Eagle Ford horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

Eagle Ford Shale:




(a)




(b)

Figure 11. Eagle Ford Shale horizontal wells' water use intensity as a function of (a) depth; and (b) formation thickness.

Eagle Ford Shale:



(a)



(b)

Note: $25 \text{ km}^2 = 154 \times 40$ acres, that is, $154 \text{ wells}/25 \text{ km}^2 = 1 \text{ well}/40 \text{ acres}$

Figure 12. Eagle Ford Shale spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).

Eagle Ford Shale:





Figure 13. Eagle Ford Shale county-level average lateral spacing.

Table 4. Eagle Ford Shale county-level average lateral spacing for top producing counties.

County Name	Sum lateral length / county area (km/km ²)	Average Lateral Spacing (1000 ft)
Karnes	0.236	13.93
Dimmit	0.162	20.30
La Salle	0.116	28.20
De Witt	0.111	29.63
Gonzales	0.080	41.01
McMullen	0.075	43.79
Webb	0.080	41.11


TX-Haynesville Shale:




Note: red squares represent average ; blue diamonds represent median; only partial data for 2012

Figure 14. TX-Haynesville Shale horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

TX-Haynesville Shale:





(a)



(b)

Figure 15. TX-Haynesville Shale horizontal water use intensity as a function of (a) depth; and (b) formation thickness.

TX-Haynesville Shale:



Note: $25 \text{ km}^2 = 154 \times 40$ acres, that is, $154 \text{ wells}/25 \text{ km}^2 = 1 \text{ well}/40$ acres

Figure 16. TX-Haynesville Shale spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).

TX-Haynesville Shale:





Figure 17. TX-Haynesville Shale county-level average lateral spacing.

Table 5. TX-Haynesville Shale county-level average lateral spacing for top producing counties.

County Name	Sum lateral length / county area (km/km ²)	Average Lateral Spacing (1000 ft)
San Augustine	0.137	23.97
Shelby	0.074	44.24
Nacogdoches	0.065	50.78
Sabine	0.061	54.11
Panola	0.046	72.03
Harrison	0.045	72.84


Permian Basin, Wolfberry Verticals:



Note: red squares represent average ; blue diamonds represent median; only partial data for 2012

Figure 18. Wolfberry verticals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median vertical productive section length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.


Permian Basin, Wolfcamp Horizontals:



Note: red squares represent average ; blue diamonds represent median; only partial data for 2012

Figure 19. Wolfcamp horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

Permian Basin, Canyon – Horizontals:



(a)


(b)




(c)



(d)



(e)




(f)

Note: red squares represent average ; blue diamonds represent median; only partial data for 2012


Figure 20. Canyon Sand horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

Permian Basin, Clearfork - Verticals




(a)

(b)



(c)

(d)




(e)

(f)

Note: red squares represent average ; blue diamonds represent median; only partial data for 2012


Figure 21. Clearfork verticals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median vertical productive section length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

Permian Basin, San Andres-Grayburg -Verticals




(a)

(b)



(c)

(d)




(e)

(f)

Note: red squares represent average ; blue diamonds represent median; only partial data for 2012


Figure 22. San Andres-Grayburg verticals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median vertical productive section length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

Permian Basin, San Andres-Grayburg -Horizontals




(a)

(b)



(c)

(d)



(e)

(f)

Note: red squares represent average ; blue diamonds represent median; only partial data for 2012

Figure 23. San Andres-Grayburg horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

Permian Basin:







Figure 24. Permian Basin spatial distribution of water intensity for (a) vertical and (b) horizontal wells.

Permian Basin:



(a)

Note: $25 \text{ km}^2 = 154 \times 40$ acres, that is, $154 \text{ wells}/25 \text{ km}^2 = 1 \text{ well}/40$ acres



(b)

Figure 25. Permian Basin spatial distribution of (a) vertical well density and (b) density of lateral (cumulative length per area) for horizontal wells.

Permian Basin






Figure 26. Permian Basin county-level average lateral spacing

Anadarko Basin: Granite Wash Horizontals:




(a)


(b)




(c)



(d)



(e)




(f)

Note: red squares represent average ; blue diamonds represent median; only partial data for 2012


Figure 27. Granite Wash horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

Anadarko Basin: Cleveland Horizontals:




(a)

(b)



(c)

(d)




(e)

(f)

Note: red squares represent average ; blue diamonds represent median; only partial data for 2012

Figure 28. Cleveland horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

Anadarko Basin: Marmaton Horizontals:



Note: red squares represent average ; blue diamonds represent median; only partial data for 2012

Figure 29. Marmaton horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

Anadarko Basin: Granite Wash Horizontals:




Figure 30. Granite Wash horizontal water use intensity as a function of depth.

Anadarko Basin: Cleveland Horizontals:






Figure 31. Cleveland horizontal water use intensity as a function of depth.

Anadarko Basin: Granite Wash Horizontals:



(a)



(b)

Note: $25 \text{ km}^2 = 154 \times 40 \text{ acres}$, that is, $154 \text{ wells}/25 \text{ km}^2 = 1 \text{ well}/40 \text{ acres}$

Figure 32. Granite Wash spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).

Anadarko Basin: Granite Wash Horizontals:






Figure 33. Granite Wash horizontals county-level average lateral spacing

Table 6. Granite Wash county-level average lateral spacing for top producing counties

County Name	Sum lateral length / county area (km/km ²)	Average Lateral Spacing (1000 ft)
Wheeler	0.351	9.34
Hemphill	0.082	39.74
Roberts	0.036	90.54


Anadarko Basin: Cleveland Horizontals:




Note: $25 \text{ km}^2 = 154 \times 40 \text{ acres}$, that is, $154 \text{ wells}/25 \text{ km}^2 = 1 \text{ well}/40 \text{ acres}$

Figure 34. Cleveland spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).

Anadarko Basin: Marmaton Horizontals:



(a)




(b)

Note: $25 \text{ km}^2 = 154 \times 40 \text{ acres}$, that is, $154 \text{ wells}/25 \text{ km}^2 = 1 \text{ well}/40 \text{ acres}$

Figure 35. Marmaton spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).

Anadarko Basin: Horizontals:



Note: $25 \text{ km}^2 = 154 \times 40 \text{ acres}$, that is, $154 \text{ wells}/25 \text{ km}^2 = 1 \text{ well}/40 \text{ acres}$




Figure 36. Anadarko spatial distribution of (a) water intensity; and (b) density of lateral (cumulative length per area).






Figure 37. Map view of wells' lateral expression and vertical well location in the Anadarko Basin.

East Texas Basin: Cotton Valley Verticals



Note: red squares represent average ; blue diamonds represent median; only partial data for 2012
 Figure 38. Cotton Valley verticals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median vertical productive section length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

East Texas Basin: Cotton Valley Horizontals



Note: red squares represent average ; blue diamonds represent median; only partial data for 2012
 Figure 39. Cotton Valley horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

East Texas Basin: Cotton Valley






Figure 40. Cotton Valley horizontal water use intensity as a function of depth.



Note: $25 \text{ km}^2 = 154 \times 40 \text{ acres}$, that is, $154 \text{ wells}/25 \text{ km}^2 = 1 \text{ well}/40 \text{ acres}$

Figure 41. Cotton Valley spatial distribution of density of lateral (cumulative length per area).

East Texas Basin: Cotton Valley




Note: $25 \text{ km}^2 = 154 \times 40$ acres, that is, $154 \text{ wells}/25 \text{ km}^2 = 1 \text{ well}/40 \text{ acres}$

Note: Cotton Valley wells drilled before 2005 are not included (see Nicot et al., 2011 for details).

Figure 42. Cotton Valley spatial distribution of density of vertical wells (years 2005-2011).

Gulf Coast Basin, Olmos - Horizontal



Note: red squares represent average ; blue diamonds represent median; only partial data for 2012
 Figure 43. Olmos horizontals, various historical parameters and coefficients for reported and estimated water use as a function of time: (a) number of wells; (b) water use; (c) average/median lateral length; (d) average/median water use per well; (e) average/median water use intensity; (f) average/median proppant loading.

III-2. Current Water Consumption and Sources

III-2-1 Information about Recycling/Reuse and Brackish Water Use

We collected information about recycling/reuse and brackish water use gathered during discussions with operators (Table 7). The amount of fresh water used is quite unequal across the different plays as a function of the local conditions. It can be as low as 20% in Far-West Texas or nearly 100% in East Texas. Collecting a sufficient amount of information concerning recycling/reuse and brackish water use is an improvement over the 2011 report which overall underestimated it. Reuse is limited by the amount of flow back that varies across plays. We could not document volumes of water recycled from wastewater treatment plants, but the TCEQ lists ~30 municipal and industrial facilities located in the Barnett Shale and Eagle Ford Shale plays that provide water to the industry (Figure 44). Groundwater/surface water could be extremely variable within a single play, but water data also reflect local conditions (Table 8): heavy surface water use towards the eastern part of the state and reliance on groundwater (sometimes brackish) elsewhere. The following short paragraphs discuss recycling/reuse and brackish water use and GS/SW split in major plays/regions.

Barnett Shale: For the most part, operators use fresh surface water in this play (estimated at 80% of “new” water). This is a change from the 50%+ groundwater use estimated in 2006 in Bené et al. (2007) and Nicot and Potter (2007). Some operators use brackish water, particularly in the combo play and on the western edges of the play. Some also use outfall from wastewater treatment plants. Overall, little recycling/reuse and brackish water use is currently occurring in this play as compared to other plays further west or south.

Eagle Ford Shale: Operators rely mostly on groundwater (estimated at 90% of “new” water) and there is a significant amount of brackish water being used (currently estimated at 20% but variable among operators). Several aquifers are brackish in the footprint of the play: the Gulf Coast aquifers and the Wilcox aquifers as well as the downdip section of the Carrizo aquifer.

Haynesville Shale and East Texas Basin: Water is generally plentiful in East Texas and no significant recycling/reuse and use for brackish water was documented during this study. We estimated it at 5%, mostly from treatment plants and produced water from Cotton Valley wells. We estimated that about 70% of the “new” water is groundwater.

Permian Basin: A significant percentage (30% or more) of the HF water used in both the Midland and Delaware basins is brackish. Nearly all of the water used is groundwater tapping aquifers such as the Ogallala (which is often brackish towards its southern domain, where the industry has many HF operations), and the Dockum, Trinity Edwards, Capitan, and other aquifers. The industry currently does little recycling/reuse, although several companies use produced water from conventional oil and gas operations. Such produced water has relatively low salinity at several places in the basin.

Anadarko Basin: This basin has hosted much recycling/reuse (estimated at 20%) and use of brackish water (estimated at 30%). Most of the “new” water is groundwater (estimated at 80%).

III-2-2 2011 HF Water Use and Consumption

Combining information collected from the IHS database, industry information, and selected information from the 2011 report results in an estimated water use for HF of ~81,500 AF across the state in 2011 (Table 9). The Barnett Shale and the Eagle Ford shale used a similar amount of

water (~25 kAF), but less fresh water was used in the Eagle Ford. The Permian Basin is catching up (~15 kAF), but it uses relatively less fresh water than the two shale plays. Water use in the Texas section of the Haynesville Shale is becoming subordinate to other plays located in the same area (for example, Cotton Valley). County-level water use (Table 10) shows that many counties across the state have some HF water use (126 counties with >1AF in 2011 and 26 counties with >1kAF). The top 10 HF users consist of Tarrant County in the Barnett core (8.8 kAF), Webb County in the southern Eagle Ford (4.6 kAF), Johnson County in the core of the Barnett Shale (4.2 kAF), Karnes County in the Eagle Ford (3.9 kAF), Wheeler County in the Granite Wash of the Anadarko Basin (3.8 kAF), Dimmit County in the Eagle Ford (3.7 kAF), Denton County in the core of the Barnett Shale (3.2 kAF), Montague County in the combo play of the Barnett Shale (3.2 kAF), La Salle County in the Eagle Ford (2.9 kAF), and Wise County in the core of the Barnett Shale (2.3 kAF). The top ten counties total about half of the HF water use in the state. The top 10 counties stay the same when only water consumption is considered despite some reshuffling because of the variable impact of recycling/reuse and brackish water use.

In the next section we compare our current findings to the findings of the 2011 report (that projected a water use of 62 kAF in 2011, Table 9) and explain the discrepancies.

Table 7. Estimated percentages of recycling/ reused and brackish water use in main HF areas in 2011.

Play / Region	Type	Current (2011) %
Permian Far West	Recycled/reused	0%
	Brackish	80%
	Fresh	20%
Permian Midland	Recycled/reused	2%
	Brackish	30%
	Fresh	68%
Anadarko Basin	Recycled/reused	20%
	Brackish	30%
	Fresh	50%
Barnett Shale	Recycled/reused	5%
	Brackish	3%
	Fresh	92%
Eagle Ford Shale	Recycled/reused	0%
	Brackish	20%
	Fresh	80%
East Texas Basin	Recycled/reused	5%
	Brackish	0%
	Fresh	95%

Table 8. Estimated groundwater / surface water split (does not include recycling / reuse)

Play / Region	Groundwater	Surface Water
Barnett Shale	20%	80%
Eagle Ford Shale	90%	10%
East Texas Basin	70%	30%
Anadarko Basin	80%	20%
Permian Basin	100%	0%

Table 9. HF water use in 2008 and 2011 compared to the 2011 projected water use from 2008.

Play / Region Unit: kAF	2011 Actual Water Use	Fraction Non-R/R Non-brackish	2011 Actual Water Consumption	2011 Projected Water Use
Barnett Shale	25.75	0.92	23.69	33.08
Eagle Ford Shale	23.76	0.8	18.81	10.07
East Texas Basin	7.54	0.95	7.06	8.46
Anadarko Basin	6.52	0.5	3.21	2.26
Permian Basin	14.44	0.68 / 0.2	8.55	7.26
Gulf Coast Basin	3.49	0.95 / 0.8	3.31	1.00
Statewide	81.51	0.79*	64.63	62.13

FrackingWaterUse2008&2011_Bob-JPComp_2.xls

*: computed from state consumption and use columns (sum of other rows)


Table 10. County-level estimate of 2011 HF water use and water consumption (kAF).

County	HF Water Use (kAF)	HF Water Consumption (kAF)	County	HF Water Use (kAF)	HF Water Consumption (kAF)
Andrews	1.391	0.946	Limestone	0.268	0.214
Angelina	0.007	0.006	Lipscomb	0.382	0.191
Archer	0.017	0.016	Live Oak	0.972	0.777
Atascosa	1.009	0.807	Loving	0.189	0.038
Bee	0.066	0.053	McMullen	1.752	1.401
Borden	0.033	0.023	Madison	0.204	0.163
Brazos	0.238	0.191	Marion	0.010	0.010
Brooks	0.008	0.006	Martin	2.035	1.384
Burleson	0.247	0.197	Maverick	0.192	0.154
Caldwell	0.075	0.060	Midland	1.573	1.070
Carson	0.085	0.042	Milam	0.034	0.027
Cherokee	0.010	0.009	Mitchell	0.018	0.012
Clay	0.058	0.053	Montague	3.221	2.963
Cochran	0.031	0.021	Moore	0.076	0.038
Coke	0.001	n/a	Nacogdoches	1.128	1.072
Cooke	1.480	1.362	Newton	0.098	0.093
Crane	0.159	0.108	Nolan	0.011	0.008
Crockett	0.475	0.323	Nueces	0.016	0.013
Crosby	0.012	0.008	Ochiltree	0.273	0.136
Culberson	0.166	0.033	Orange	0.006	n/a
Dallas	0.079	0.073	Palo Pinto	0.041	0.038
Dawson	0.089	0.061	Panola	0.966	0.917
Denton	3.249	2.989	Parker	1.086	1.000
DeWitt	2.151	1.721	Pecos	0.110	0.022
Dimmit	3.706	2.965	Polk	0.133	0.126
Ector	0.756	0.514	Potter	0.044	0.022
Ellis	0.038	0.035	Reagan	1.240	0.843
Erath	0.012	0.011	Reeves	0.522	0.104
Fayette	0.132	0.106	Roberts	0.393	0.197
Franklin	0.014	0.014	Robertson	0.306	0.245
Freestone	0.424	0.339	Runnels	0.004	0.003
Frio	0.729	0.583	Rusk	0.158	0.150
Gaines	0.142	0.096	Sabine	0.147	0.139
Garza	0.001	n/a	San Augustine	1.622	1.541
Glasscock	1.434	0.975	Schleicher	0.090	0.061
Gonzales	2.224	1.779	Scurry	0.010	0.007
Grayson	0.021	0.020	Shackelford	0.002	0.002
Gregg	0.025	0.024	Shelby	1.419	1.348
Grimes	0.095	0.076	Sherman	0.002	0.001
Guadalupe	0.018	0.014	Smith	0.005	0.005
Hansford	0.011	0.005	Somervell	0.287	0.264
Hardeman	0.017	0.012	Starr	0.036	0.029
Harrison	0.893	0.849	Sterling	0.057	0.039
Hemphill	1.462	0.731	Stonewall	0.001	n/a
Henderson	0.012	0.012	Sutton	0.034	0.023
Hidalgo	0.059	0.047	Tarrant	8.805	8.101
Hill	0.131	0.120	Terrell	0.010	0.007
Hockley	0.005	0.003	Terry	0.003	0.002
Hood	0.645	0.593	Titus	0.003	0.003

County	HF Water Use (kAF)	HF Water Consumption (kAF)	County	HF Water Use (kAF)	HF Water Consumption (kAF)
Houston	0.178	0.142	Tyler	0.076	0.072
Howard	0.552	0.376	Upshur	0.004	0.004
Hutchinson	0.005	0.002	Upton	1.761	1.198
Irion	0.875	0.595	Ward	0.568	0.114
Jack	0.048	0.044	Washington	0.036	0.029
Jasper	0.087	0.083	Webb	4.596	3.677
Johnson	4.192	3.857	Wheeler	3.792	1.896
Karnes	3.869	3.095	Wilson	0.417	0.334
Kenedy	0.006	0.005	Winkler	0.062	0.012
Kleberg	0.034	0.028	Wise	2.314	2.129
La Salle	2.901	2.321	Yoakum	0.018	0.013
Lavaca	0.118	0.094	Young	0.008	0.007
Lee	0.131	0.105	Zapata	0.032	0.026
Leon	0.273	0.218	Zavala	0.407	0.127
			SUM	81.50 kAF	64.63 kAF

Note: filtered at 0.001 kAF

FrackingWaterUse2008&2011_Bob-JPComp_2.xls



Source: TCEQ, 2012

Figure 44. Location of waste water treatment facilities that provide or have provided water to the industry for HF as of July 2012.

III-3. Comparison to Earlier Findings

Projections made in 2009 for 2011 in the 2011 report underestimated water use by about 30% (81.5 kAF compared to 62.1 kAF, Table 9). It is important to understand the underlying causes in order to develop better projections in this document. Comparing actual water use in 2008 and 2011 (Figure 45) shows (1) extension of HF across the state, Barnett Shale stays relatively steady, fracturing in the Haynesville Shale and Anadarko Basin expands, and the Eagle Ford becomes much more prominent as does the Permian Basin. A bar plot illustrates the county-by-county discrepancies between projections and actual numbers (Figure 46). A cross-plot is a different way of presenting the same information (Figure 47), and it is apparent that most counties with larger water use (dots in the upper right-hand side of the side) were correctly accounted (no dots on either the x- or y-axis), even if it was underestimated (dots mostly below the 1:1 line). Major discrepancies occurred because there was no Barnett extension outside of the core area (for example, Bosque, Comanche, Erath, and Palo Pinto counties in Figure 46), and because of more and faster development in the Eagle Ford Shale and Permian Basin. Both these factors are connected to the drop in gas price and increase in oil price in the past 2 or 3 years, parameters notoriously difficult to predict.




Figure 45. Spatial distribution of HF water use in 2008 and 2011.





Figure 46. Bar plot comparison of 2011 actual water use to projections from 2009.



Note: Note the log-log scale.

Figure 47. County-level cross-plot comparison of 2011 actual water use to projections from 2008. Values on x- and y- axis represent counties whose actual (y-axis) / projected (x-axis) water use is 0. A total of 168 counties are represented.

III-4. Drilling Water Use

In the course of the study, we also collected information about drilling water use. Results are not sufficiently representative to change results presented in the 2011 report amounting to 8 kAF. The general observation, though, is that drilling requires water of better quality than HF although in smaller amounts (Table 11). The amount of water used depends on the length of the well and on operator preferences but also, more importantly, heavily on local factors. For example, in the Eagle Ford the drilling muds used in drilling through horizontal sections (for example, Fan et al., 2011) are oil-based.

Table 11. Drilling water use information

Play / Region in 1000's gal/well	Range provided by operators	Comments
Barnett Shale	250	N/A
	210-420	~Fresh
	168	~Fresh
	500	~Fresh
Eagle Ford Shale	125	N/A
	420	N/A
	160	~Fresh
	126	~Fresh
	252-420	~Fresh
East Texas Basin	600	N/A
	840-1,100	~Fresh
	420	~Fresh
Anadarko Basin	200	N/A
	420	~Fresh
Midland Basin (Permian Basin)	84	~Fresh
	100	N/A
	210	~Fresh
	210-420	~Fresh
Delaware Basin (Permian Basin)	100	N/A
	210-420	Brackish

Note: fresh is defined as TDS < 3,000 mg/L

IV. Water Use Projections

This section describes projections for HF water use and fresh-water consumption in Texas to year 2060. As described in the 2011 report, all projections entail many uncertainties and those caveats are still valid in this update. In general, the life of the plays was extended beyond 2060, less prospectivity was given to the gas window, and steeper development to the oil window section of plays or tight oil plays. The overall results is that the HF water use will have a broad plateau at ~125 kAF/yr around the 2020-2030 decade and then slowly decrease with time to 2060 and beyond (Figure 48). However, the amount of fresh water consumed (that is, not recycled or reused or brackish water) will stay relatively constant at ~70 kAF despite the increase in water use and then slowly subside with the decrease in HF activities. Fresh-water use will decrease for two reasons: (1) the industry is getting better at reusing flow back (but sometimes limited by the small fraction coming back) and at finding alternate sources of recycling (treatment plants, produced water from conventional wells) and at using brackish water because of the technological advances in additives tolerating more saline water. And (2) the Permian Basin, which may become the focus of HF in Texas in the long run, offers great production potential. In the Permian Basin, fresh water is at a premium and brackish water is already used by the industry.

Total oil and gas water use and consumption (combining HF, waterflooding, and drilling) is presented in Figure 49. Oil and gas water use, consistent with the definition of make-up fresh water used in this document, was computed by summing HF water use (Figure 48), drilling water use –with no change from the 2011 report, and waterflood water use –computed from the 2011 report by adding fresh and brackish water use. Oil and gas water consumption was computed by summing HF water consumption (Figure 48), drilling water use –with no change from the 2011 report and the additional note that water use and consumption are identical. Waterflood water consumption is the same as water use in the 2011 report that represented fresh water use. Projected oil and gas water use and consumption are dominated by HF. By design, in the 2011 report, drilling technology was projected to move the industry away from the use of fresh water. Progress in waterflooding was also projected to decrease fresh water requirements but to increase brackish water use until the whole industry relies only on saline water (not showed). Under these assumptions, oil and gas industry water use is projected to peak with a broad plateau at 180 kAF in the 2020-2030 decade, slowly declining to ~60 kAF by 2060. Fresh water consumption in the oil and gas industry is projected to reach a maximum of ~100 kAF before the end of this decade and then to slowly decrease to a low level of a few tens of thousands AF by the middle of the century.

We did not account for many unknowns that could possibly impact the results as they did in the Eagle Ford Shale when the industry switched from slick-water fracs to gel fracs in the oil window that use less water. The Eagle Ford was the only play in which we observed such a trend, everywhere else the trend (based on 2 to 5 years of data) shows an increase or a steady value in water intensity (Table 12). Data about recycling/reuse and brackish water use were derived from industry information of these uses as of today and in 2020 (Table 13). The most likely values from 2011 and 2020 are essentially estimated directly from the various responses in a given play. Extrapolation to 2060 and translation to high and low scenarios for all years starting in 2012 are speculative and are based on industry trends and on the general knowledge of the authors about fresh and brackish water aquifers and of their yields around the state. The

amount of reuse cannot be larger than the amount of flow back / produced water from recently fractured wells and at the play level reuse is likely less because of the operational issues of transporting water. Some plays, such as the Haynesville and Eagle Ford Shales, are at a disadvantage for this; they produce back less than 20% of the injected water (Table 14). They, and others, could however take advantage of produced water from other formations.

We did not deviate much from the overall water use of the 2011 report because of constraints accounted for the 2011 report and related to drilling rig count, labor force availability/staff shortage, infrastructure development, and other factors. National rig count seems steady at ~2,000 or slightly lower in the past year (~50% of them in Texas), but drillers are improving at operating them, which suggests that the projections presented in this update are consistent with the number of drilling rigs currently available.

Cumulative water use is related to the eventual well density or lateral spacing. Ultimate average spacing between laterals, or vertical well density, is the parameter driving water use along with water intensity. Typical vertical well spacing is 1 well per 40 acres; that ratio can decrease to 1 well per 20 or 10 acres in some instances. Typical lateral spacing can be computed from 1 horizontal well per 160 acres. If lateral length is 5,000 ft, the resulting spacing between laterals is 1,400 ft. If the horizontal well density declines to 1 well per 40 acres, lateral spacing is 350 ft. This update document assumes a lateral spacing of 1000 ft, perhaps smaller in oil windows (Figure 51).

County-level projections for HF water use and water consumption are listed in Table 15. The county coverage is essentially the same as in the 2011 report with the addition of four counties in East Texas (Polk, Tyler, Jasper, and Newton counties, Figure 50). Total oil and gas (combining HF, waterflooding, and drilling) county-level projections are presented in Table 16.

The following paragraphs address HF projection issues specific to each play and region. Each play is represented by two plots. One plot compares projections from the 2011 report to projections from this update. The second plot displays water use and fresh water consumption in the high, low, and most likely scenarios. Only the latter is displayed in the first plot and is retained as the preferred set of projections to be used by the TWDB. As explained in the Methodology Section (Section II), low and high scenarios were derived by varying two factors: (1) the prospectivity factor, which assesses the ultimate amount of HF in a play, varies on a county and play basis from 1 to 0, with 1 meaning the county is within the core area and highly prospective (for example, Tarrant County in the Barnett Shale) and near- zero values suggesting that little of the county will be developed (for example, Shackelford County in the Barnett Shale); and (2) coefficients for recycling/reuse and brackish water use (Table 13). The prospectivity factor was changed according to a sliding linear scale: a value of 1 stays at 1 but a value of 0.2 either goes to zero (low water use scenario) or 0.4 (high water use scenario). The change was made systematically with no tentative exercise to tailor it to each county/play couple. In the case of tight oil/ tight gas plays, a third factor was varied. This factor varies from 0 to 1 and addresses the spatial coverage of the county that could ultimately undergo HF. In the case of resource plays such as shale plays, the factor is constant and close to one because the whole footprint of the play is potentially a target for drilling. The only unknown is the well density which is accounted for through the prospectivity factor. In tight oil/gas plays, it cannot be assumed that the whole footprint of the formation will experience HF because some parts of it can be properly produced through conventional wells. This third factor was used in the East

Texas (Cotton Valley), Anadarko (Granite Wash), Gulf Coast (Austin Chalk), and Permian basins.

Barnett Shale: In this play with the longest history, we considerably decreased the prospectivity factors outside of the core area in the most likely scenario. That is, instead of increasing water use because of the expansion of the productive Barnett Shale footprint, we assumed that most of the HF will stay confined to the core area and stay relatively stable for a few years before slowly decreasing (Figure 52a). The peak from earlier projections has disappeared and water use should stay below 30 kAF and decrease more slowly than projected in the 2011 report. The high water use scenario projection (Figure 52b) displays a small increase in water use (but not in water consumption) in the 2020 decade because the prospectivity factors are closer to those used in the 2011 report.

Eagle Ford Shale: Projections for this play display a decrease in water use compared to those projected values of the 2011 report (Figure 53a) because of the observed decrease in water intensity that we assumed will hold in the future. The projections suggest a slow increase in water for the next 10 years with a broad peak at ~35kAF and a slow decrease beyond 2060. Unlike the Barnett with a clearly delimited core, we assumed that most counties in the Eagle Ford are highly prospective and thus there is not much variation between high and low scenario projections except when recycling/reuse and use of brackish water are included (Figure 53b).

Pearsall Shale: This gas play was briefly hydraulically fractured in the mid-2000's and has not received a lot of attention since then. However, initial production estimates suggest that the play will be produced in the future. We used the same water use parameters in the Pearsall as those in the Eagle Ford Shale because these plays are geographically close. Projections from the 2011 report were only slightly modified displacing the peak water use at ~10 kAF by about 5 years into the future (Figure 54a). As was the case for the Eagle Ford, the high and low scenarios are mostly impacted by the amount of recycling/reuse and brackish water use (Figure 54b).

TX-Haynesville and Bossier Shales: The Haynesville and Bossier Shales have declined in operator interest because of their relatively high operational cost and low gas prices. They are, however, still likely to produce significant amounts of gas in the future, albeit at a lower rate than anticipated in the 2011 report. Projections of this update document show a decreased and broader peak (Figure 55a), with annual water use slated to be no higher than ~12kAF. A minor player, the Haynesville-West play will possibly undergo some development on the western flank of the East Texas Basin and its water use projections stay similar to that of the 2011 report (Figure 56a), with a decrease peak as well. Low and high scenario projections stay relatively close together (Figure 55b), because there is little variability in terms of projected non-fresh water use (almost none).

Other East Texas Formations: This category includes all formations except the Haynesville and Bossier Shales, such as the Cotton Valley, James Lime, Bossier Sands, and others. The same water consumption data used in the Haynesville were used for this group of formations. Relative to the 2011 report projections, the projections derived in this update assumed a broader peak displaced toward the future by ~10 years (Figure 57a). Projected maximum water use is estimated at <5 kAF/yr. The small variance between water use and water consumption is explained by the location of the plays in East Texas where fresh water is relatively abundant and the large differences between the different scenario projections is due to the spread of the third factor, addressing spatial coverage of the formation of interest (Figure 57b).

Gulf Coast Formations: Amount of water use and consumption in the Gulf Coast Basin outside of the shale plays is very uncertain. The Gulf Coast Basin is the area in Texas that has experienced the least HF (Nicot et al., 2011) and explained the large range of projections between the different scenarios (Figure 58b). This category include formations such as the Olmos Sands and the Austin Chalk, and these projections assumed that water use will peak at ~8kAF in the 2020's (Figure 58a). Water consumption is assumed to be much lower because most of the plays are in South Texas, where there are some brackish water resources.

Anadarko Basin: Anadarko Basin consists mostly of the Granite Wash in Hemphill and Wheeler counties and the Marmaton/Cleveland in Ochiltree and Lipscomb counties. Current water use in this basin is much higher than anticipated in the 2011 report projections. We revisited prospectivity factors and the projected water use reaches a broad peak of ~9kAF in the 2020's (Figure 59a) with a smaller projected water consumption because of anticipated recycling/reuse and brackish water use. However, the uncertainty in final coverage put this basin in the same category as the Gulf Coast Basin and East Basin category, resulting in a large spread of potential outcomes (Figure 59b).

Permian Basin: As has the Anadarko Basin, the Permian Basin has grown much faster than anticipated and water use projections call for a plateau at ~40 kAF during the 2020-2040 period (Figure 60a) concomitant with a fairly stable fresh water consumption at 10-15 kAF. The large gap between water use and water consumption, much larger than presented in the 2011 report (Figure 60a), is due to the expectation of availability of significant amounts of brackish water and of their extensive use by the industry (as currently documented by anecdotal evidence). The large range in outcome from the different scenarios is related to the unknowns in spatial coverage of the non-shale plays (Figure 60b). We now turn to the description of the major components making up water use in the Permian Basin. Although the Barnett-Woodford system in the Permian Basin has received limited interest, we assume it will produce gas in the future (Figure 61a). The most likely scenario calls for a peak at ~5 kAF in 2035 but with the possibility of a high scenario with a much higher water use and a low scenario with no development. Development centered on the Wolfcamp is more certain and differences between high and low scenario projections were derived mostly from assumptions on the level of use of non-fresh water (Figure 61b). The other formations in the Permian Basin also display the same uncertainty related to the amount of spatial coverage ("third factor" as described above). The most likely scenario projection is estimated to have a broad peak in the 15-20 kAF range for many years with considerably less water consumption (Figure 61c).

Table 12. Recent trends in well completion and water use in hydraulic-fractured plays.

Play	Well Type	~# of Recent Wells/yr	Recent Trend (well/yr)	Water Use / well (Mgal)	Water Use Intensity (gal/ft)	Recent Trend (water use)
Barnett	H	1500	down / steady	n/a	1200	steady
Eagle Ford	H	1000	strongly up	n/a	850	down
TX-Haynesville	H	250	up	n/a	1400	steady
Granite Wash	H	250	strongly up	n/a	1200	steady / up
	V	60	strongly down	1500	800	steady
Cleveland	H	100	steady	n/a	250	steady
	V	20	down	1.7	2000	steady
Marmaton	H	30	strongly up	n/a	250	steady
	V	10	steady	1.0	2500	up
Cotton Valley	H	100	up	n/a	1000	steady
	V	300	strongly down	0.8	1200	steady
Olmos	H	50	up	n/a	1000	up
	V	100	strongly down	0.15	2500	steady
Wolfcamp	H	150	strongly up	n/a	900	strongly up
Wolfberry	V	2000	up	1.0	350	up
Canyon	V	300	down	0.4	500	up
Clear Fork	V	800	up	0.8	350	up
San Andres	H	50	strongly down	n/a	350	strongly up
	V	800	steady / up	0.15	500	steady

Table 13. Coefficients (%) to compute water consumption to be applied to total water use.

Play / Region		High Water Use	Most Likely	Low Water Use
Far West Permian Basin	Recycling			
	2011	0	0	0
	2020	0	50	40
	2060	0	40	40
	Brackish			
	2011	80	80	80
	2020	80	30	50
2060	80	40	50	
Permian Midland Basin	Recycling			
	2011	2	2	2
	2020	2	25	30
	2060	2	30	40
	Brackish			
	2011	30	30	30
	2020	30	40	40
2060	30	40	50	
Anadarko Basin	Recycling			
	2011	20	20	20
	2020	20	30	40
	2060	20	40	40
	Brackish			

Play / Region		High Water Use	Most Likely	Low Water Use
	2011	30	30	30
	2020	30	30	30
	2060	30	30	40
Barnett Shale	Recycling			
	2011	5	5	5
	2020	5	10	25
	2060	5	20	20
	Brackish			
	2011	3	3	3
	2020	3	15	20
2060	3	25	25	
Eagle Ford Shale	Recycling			
	2011	0	0	0
	2020	0	10	10
	2060	0	10	10
	Brackish			
	2011	20	20	20
	2020	20	40	50
2060	20	50	50	
South Texas	Recycling			
	2011	0	0	0
	2020	0	10	10
	2060	0	10	10
	Brackish			
	2011	20	20	20
	2020	20	40	50
2060	20	50	50	
East Texas	Recycling			
	2011	5	5	5
	2020	5	10	10
	2060	5	10	10
	Brackish			
	2011	0	0	0
	2020	0	0	10
2060	0	10	10	

Table 14. Estimated flow back/produced water volume relative to HF injected volume.

Play / Region	Comment
Delaware Basin (Permian Basin)	Close to 100% in year 1, 150% well life >200% well life
Midland Basin (Permian Basin)	50%-100% in year 1
Anadarko Basin	~50% in month 1, 90% at month 6
Barnett Shale	10-20% month 1, 20-60% well life 70% year1; 150% in 5 years
Eagle Ford Shale	20% over life; 20% over life
Haynesville Shale	20% over life; 15% over life
Cotton Valley Fm.	60% month 1, >100% well life; 40% or 100% over life

Table 15. County-level estimate of 2012-2060 projections for HF water use and water consumption (AF).

County	Water Use (AF)										Water Consumption (AF)											
	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060
Anderson	0	31	58	89	119	131	139	124	105	85	66	0	23	41	64	86	97	104	92	76	61	46
Andrews	1,391	1,617	2,140	2,053	1,965	1,878	1,654	1,431	1,207	983	806	946	862	749	690	634	580	501	425	351	279	224
Angelina	7	60	160	260	360	379	345	310	276	241	207	6	56	144	231	315	327	293	260	228	196	165
Aransas	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Archer	17	81	183	284	385	354	321	289	257	225	193	16	68	137	206	270	239	209	181	154	129	106
Armstrong	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Atascosa	1,009	2,902	2,638	2,589	2,594	2,598	2,602	2,314	1,953	1,591	1,230	807	2,064	1,583	1,500	1,443	1,386	1,329	1,144	935	736	545
Austin	0	0	98	195	293	264	234	205	176	146	117	0	0	59	115	169	148	129	110	92	75	59
Bailey	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bandera	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Basin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Baylor	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bee	66	80	101	108	94	81	67	54	40	27	13	53	60	64	67	57	48	39	31	23	15	7
Bell	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bexar	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Blanco	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Borden	33	228	638	892	899	906	764	622	480	338	230	23	122	223	307	303	300	248	198	150	104	69
Bosque	0	192	329	466	603	553	502	462	402	352	301	0	162	247	338	422	373	327	283	241	202	166
Bowie	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Brazoria	0	41	60	79	97	91	79	67	55	43	31	0	31	38	49	59	54	46	38	31	24	17
Brazos	238	322	696	931	1,166	1,036	905	775	644	514	384	191	243	431	559	681	592	506	423	343	267	193
Brewster	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Briscoe	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Brooks	8	37	49	62	62	54	46	38	30	22	14	6	28	31	38	38	32	27	22	17	12	8
Brown	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Burleson	247	331	943	1,409	1,877	1,676	1,474	1,273	1,071	867	665	197	250	580	840	1,090	952	819	690	567	447	334
Burnet	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Calwell	75	90	116	103	90	77	64	52	39	26	13	60	68	73	64	55	46	38	29	22	14	7
Callahan	0	25	33	42	42	37	31	26	21	15	10	0	19	21	26	26	22	18	15	11	8	5
Callahan	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cameron	0	37	50	62	62	54	46	38	30	22	14	0	28	31	38	38	32	27	22	17	12	8
Camp	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Carson	85	0	0	0	0	0	0	0	0	0	0	42	0	0	0	0	0	0	0	0	0	0
Cass	0	10	25	41	56	68	60	52	45	37	30	0	9	24	38	52	60	52	45	38	31	24
Castro	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Chambers	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cherokee	10	70	128	186	244	284	253	221	190	158	126	9	66	122	173	223	254	221	189	159	129	101
Childress	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Clay	58	194	355	516	678	621	565	508	452	395	339	53	164	266	374	474	419	367	318	271	227	186
Cochran	31	94	121	149	176	203	180	158	135	113	90	21	50	42	51	59	67	59	50	42	35	27
Coke	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Coleman	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Collin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Collingsworth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Colorado	0	38	517	996	1,462	1,314	1,166	1,018	870	722	574	0	29	312	587	843	741	643	548	458	371	287
Comal	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Comanche	0	125	228	332	436	392	349	305	261	218	174	0	105	171	241	305	265	227	191	157	125	96
Concho	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cooke	1,480	1,653	1,294	934	575	215	0	0	0	0	0	1,362	1,396	970	677	402	145	0	0	0	0	0
Coryell	0	289	1,012	947	684	421	158	0	0	0	0	0	244	759	686	479	284	103	0	0	0	0

County	Water Use (AF)										Water Consumption (AF)											
	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060
Cottle	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Crane	159	339	438	559	681	802	729	656	583	510	438	108	181	153	189	223	257	229	203	177	152	128
Crockett	475	996	1,636	1,946	1,760	1,475	1,190	905	620	335	149	323	531	573	669	594	489	387	288	194	103	45
Crosby	12	0	0	0	0	0	0	0	0	0	0	8	0	0	0	0	0	0	0	0	0	0
Culberson	166	141	188	576	963	1,280	1,163	1,047	931	814	698	33	75	66	149	231	290	262	235	207	180	154
Dallam	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Dallas	79	654	1,018	848	679	509	339	170	0	0	0	73	553	763	615	475	343	220	106	0	0	0
Dawson	89	476	724	918	954	990	844	699	553	408	294	61	254	253	308	308	308	257	208	160	115	80
Deaf Smith	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Della	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Denton	3,249	3,159	2,106	1,053	0	0	0	0	0	0	0	2,989	2,667	1,579	763	0	0	0	0	0	0	0
DeWitt	2,151	1,977	1,773	1,569	1,354	1,130	907	684	460	237	14	1,721	1,407	1,065	924	780	638	500	369	243	122	7
Dickens	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Dimmit	3,706	4,777	4,765	4,857	4,871	4,834	4,232	3,489	2,746	2,002	1,259	2,965	3,407	2,828	2,774	2,669	2,534	2,145	1,710	1,294	895	516
Donley	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Duval	0	70	94	117	118	103	87	72	57	42	27	0	53	59	73	72	61	51	41	32	23	14
Eastland	0	0	424	642	550	458	367	275	184	92	0	0	0	318	465	385	309	238	172	110	53	0
Ector	756	983	1,340	1,434	1,529	1,484	1,309	1,134	959	784	644	514	524	469	478	488	451	390	332	274	219	176
Edwards	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Ellis	38	87	126	166	206	185	164	144	123	103	82	35	74	95	120	144	125	107	90	74	59	45
El Paso	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Erath	12	163	253	343	433	397	361	325	289	253	217	11	137	190	249	303	268	235	203	173	145	119
Falls	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Fannin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Fayette	132	1,081	2,329	2,093	1,822	1,526	1,229	932	636	340	43	106	773	1,402	1,236	1,054	864	681	505	337	176	23
Fisher	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Floyd	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Foard	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Fort Bend	0	35	46	58	58	51	43	36	28	21	14	0	26	29	36	35	30	25	20	16	11	7
Franklin	14	0	0	0	0	0	0	0	0	0	0	14	0	0	0	0	0	0	0	0	0	0
Freestone	424	750	975	1,229	1,424	1,404	1,241	1,076	912	748	584	339	678	846	1,042	1,196	1,164	1,012	863	720	582	449
Frio	729	1,119	1,146	1,176	1,189	1,159	1,127	1,097	947	769	589	583	809	701	708	692	647	602	559	465	364	266
Gaines	142	830	1,273	1,709	1,881	1,841	1,582	1,323	1,064	805	599	96	443	445	563	588	542	456	372	290	212	152
Galveston	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Garza	1	237	315	394	473	426	379	331	284	237	189	0	126	110	136	160	141	123	106	89	72	57
Gillespie	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Glasscock	1,434	1,938	2,621	2,466	2,311	1,978	1,646	1,313	980	648	427	975	1,033	917	848	780	655	535	419	306	198	128
Goliad	0	34	45	56	56	49	42	35	27	20	13	0	25	28	35	34	29	24	20	15	11	7
Gonzales	2,224	1,746	1,552	1,358	1,164	970	776	582	388	194	0	1,779	1,241	931	798	669	545	427	313	204	99	0
Gray	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Grayson	21	0	0	0	0	0	0	0	0	0	0	20	0	0	0	0	0	0	0	0	0	0
Gregg	25	134	224	313	402	449	405	362	318	274	230	24	127	208	284	357	391	347	305	263	223	184
Grimes	95	125	287	448	569	506	443	360	317	254	191	76	94	178	270	334	291	249	209	170	133	97
Guadalupe	18	0	0	0	0	0	0	0	0	0	0	14	0	0	0	0	0	0	0	0	0	0
Hale	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hall	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hamilton	0	251	304	253	203	152	101	51	0	0	0	0	212	228	184	142	103	66	32	0	0	0
Hansford	11	0	513	1,025	879	732	586	439	293	146	0	5	0	205	397	329	265	205	148	95	46	0
Hardeman	17	0	0	0	0	0	0	0	0	0	0	12	0	0	0	0	0	0	0	0	0	0
Hardin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Harris	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Harrison	893	1,578	2,223	2,012	1,851	1,689	1,527	1,365	1,203	1,041	880	849	1,479	2,030	1,808	1,636	1,469	1,307	1,149	996	847	704

County	Water Use (AF)										Water Consumption (AF)											
	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060
Hartley	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Haskell	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hays	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hemphill	1,462	2,484	2,231	1,978	1,724	1,470	1,217	963	710	456	203	731	1,132	892	646	533	426	325	231	143	61	
Henderson	12	46	124	201	278	333	296	259	222	185	148	12	44	117	187	254	297	259	222	186	118	
Hidalgo	59	63	83	104	105	91	78	64	51	37	24	47	47	53	64	54	45	37	28	20	13	
Hill	131	1,429	1,225	1,021	816	612	408	204	0	0	0	120	1,207	919	571	413	265	128	0	0	0	
Hockley	5	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0	
Hood	645	409	580	751	921	829	737	645	553	461	369	593	346	435	645	560	479	403	332	265	203	
Hopkins	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Houston	178	237	305	271	237	203	170	135	102	68	34	142	179	193	144	121	99	77	57	37	18	
Howard	552	1,471	2,360	2,822	2,642	2,250	1,859	1,468	1,076	685	422	376	784	826	892	745	604	468	336	210	126	
Hudspeth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Hunt	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Hutchinson	5	0	90	180	154	128	103	77	51	26	2	2	0	36	58	47	36	26	17	8	0	
Irion	875	1,478	2,429	2,889	2,613	2,190	1,766	1,343	920	497	221	595	788	850	882	725	574	428	287	152	66	
Jack	48	242	363	485	605	545	485	424	363	303	242	44	204	273	424	368	315	265	218	174	133	
Jackson	0	34	45	56	56	49	42	35	28	20	13	0	25	29	34	29	25	20	15	11	7	
Jasper	87	105	135	120	105	90	75	60	45	30	15	83	79	86	64	54	44	34	25	16	8	
Jeff Davis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Jefferson	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Jim Hogg	0	45	60	75	75	65	56	46	37	27	17	0	34	38	46	39	32	26	20	15	9	
Jim Wells	0	34	45	57	57	50	42	35	28	21	13	0	26	29	35	30	25	20	15	11	7	
Johnson	4,192	4,038	3,365	2,692	2,019	1,346	673	0	0	0	0	3,857	3,410	2,524	1,413	909	437	0	0	0	0	
Jones	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Karnes	3,869	2,749	2,457	2,165	1,863	1,554	1,245	937	629	320	11	3,095	1,956	1,475	1,073	876	686	505	331	165	6	
Kaufman	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Kendall	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Kendall	6	57	76	95	95	83	71	58	46	34	22	5	43	48	58	49	41	33	26	19	12	
Kent	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Kerr	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Kimble	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
King	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Kirney	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Kleberg	34	37	49	62	62	54	46	38	30	22	14	28	28	31	38	32	27	22	17	12	8	
Knox	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Lamar	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Lamb	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Lampasas	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
La Salle	2,901	4,432	4,425	4,532	4,621	4,698	4,147	3,440	2,732	2,025	1,318	2,321	3,154	2,612	2,563	2,499	2,427	1,659	1,265	889	530	
Lavaca	118	913	1,522	1,388	1,241	1,086	930	775	620	464	309	94	651	915	716	613	513	418	326	239	155	
Lee	131	203	392	508	624	553	484	414	345	274	204	105	152	243	305	365	316	270	184	142	103	
Leon	273	663	1,289	1,800	2,309	2,192	1,934	1,674	1,416	1,155	898	218	487	831	1,166	1,487	1,415	1,041	864	693	529	
Liberty	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Limestone	268	307	347	388	410	376	332	287	242	197	163	214	281	307	333	346	312	270	190	153	116	
Lipscomb	382	560	1,026	876	725	574	423	272	121	0	0	191	255	410	339	272	208	148	92	39	0	
Live Oak	972	783	729	676	692	720	748	776	689	575	461	777	558	439	399	392	388	364	324	261	200	
Llano	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Loving	189	313	418	561	704	690	627	565	502	439	376	38	167	146	187	227	213	169	147	127	107	
Lubbock	0	0	0	51	103	154	140	126	112	98	84	0	0	0	10	21	31	28	25	22	17	
Lynn	0	0	246	336	427	517	460	402	345	287	230	0	0	86	116	144	171	149	128	108	69	
McCulloch	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

County	Water Use (AF)										Water Consumption (AF)											
	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060
McLennan	0	53	120	187	253	228	203	177	152	127	101	0	45	90	135	177	154	132	111	91	73	56
McMullen	1,752	2,545	2,762	3,067	3,329	3,562	3,306	2,930	2,553	2,177	1,801	1,401	1,815	1,627	1,729	1,797	1,840	1,658	1,430	1,211	1,001	801
Madison	204	261	561	750	940	832	727	622	518	413	308	163	197	348	451	549	475	406	339	275	214	155
Marion	10	121	270	420	569	579	522	466	408	351	295	10	114	249	380	506	506	449	393	339	286	236
Martin	2,035	2,446	3,071	2,824	2,577	2,267	1,892	1,516	1,141	765	512	1,384	1,305	1,075	963	855	731	597	468	344	224	145
Mason	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Matagorda	0	46	61	77	77	67	57	47	37	28	18	0	35	39	48	40	33	27	27	21	15	9
Maverick	192	1,574	1,857	2,241	2,626	3,010	2,843	2,538	2,234	1,928	1,623	154	1,119	1,074	1,226	1,368	1,501	1,376	1,195	1,022	856	698
Medina	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Menard	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Midland	1,573	2,640	3,265	3,034	2,803	2,465	2,045	1,625	1,205	785	488	1,070	1,408	1,143	1,034	928	791	643	499	361	227	136
Milam	34	0	0	0	0	0	0	0	0	0	0	27	0	0	0	0	0	0	0	0	0	0
Mills	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Mitchell	18	238	317	397	476	428	381	333	286	238	190	12	127	111	136	161	142	124	106	89	73	57
Montague	3,221	3,496	2,997	2,497	1,998	1,498	999	499	0	0	0	2,963	2,952	2,248	1,810	1,398	1,011	649	312	0	0	0
Montgomery	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Moore	76	0	0	0	0	0	0	0	0	0	0	38	0	0	0	0	0	0	0	0	0	0
Morris	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Motley	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Motley	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nacogdoches	1,128	1,424	2,066	1,937	1,809	1,659	1,503	1,347	1,191	1,036	880	1,072	1,327	1,873	1,731	1,593	1,438	1,283	1,132	985	842	704
Navarro	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Newton	98	125	161	143	125	108	89	71	54	36	18	93	94	102	89	76	64	52	41	30	20	9
Nolan	11	0	0	0	0	0	0	0	0	0	0	8	0	0	0	0	0	0	0	0	0	0
Nueces	16	34	45	56	56	49	42	35	28	20	13	13	25	29	35	34	29	25	20	15	11	7
Ochiltree	273	408	748	985	815	646	476	306	136	0	0	136	186	299	382	306	234	166	103	44	0	0
Oldham	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Orange	6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Palo Pinto	41	194	356	518	680	612	544	476	408	340	272	38	164	267	376	476	413	354	298	245	196	150
Panola	966	1,412	1,988	1,801	1,655	1,511	1,366	1,221	1,077	932	787	917	1,323	1,816	1,618	1,464	1,314	1,169	1,028	891	758	630
Parmer	1,086	925	1,255	1,585	1,916	1,724	1,533	1,341	1,149	958	766	1,000	781	941	1,149	1,341	1,164	996	838	690	551	421
Parmer	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Pecos	110	130	173	387	601	746	674	601	528	456	383	22	69	60	108	156	180	161	142	123	105	87
Polk	133	180	232	206	180	155	129	103	77	52	26	126	136	147	128	110	92	75	59	43	28	14
Potter	44	0	0	0	0	0	0	0	0	0	0	22	0	0	0	0	0	0	0	0	0	0
Presidio	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Rains	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Randall	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Reagan	1,240	3,207	4,019	3,627	3,236	2,844	2,332	1,820	1,308	796	444	843	1,710	1,407	1,247	1,092	942	758	580	409	244	133
Real	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Red River	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Reeves	522	866	1,155	1,744	2,333	2,509	2,304	2,098	1,893	1,687	1,481	104	462	404	556	705	713	646	581	518	456	395
Refugio	0	32	42	53	53	46	39	33	26	19	12	0	24	27	33	32	27	23	19	14	10	7
Roberts	393	1,628	1,419	1,210	1,002	793	584	376	167	0	0	197	742	568	469	376	287	205	127	54	0	0
Robertson	306	587	741	773	806	734	639	544	449	354	259	245	501	587	619	648	584	500	419	342	268	196
Rockwall	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Runnels	4	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0
Rusk	158	477	930	1,384	1,838	1,707	1,542	1,378	1,213	1,048	884	150	446	850	1,245	1,627	1,487	1,322	1,161	1,005	853	707
Sabine	147	235	470	705	940	861	783	705	627	548	470	139	218	423	625	823	743	666	590	517	445	376
San Augustine	1,622	2,092	1,953	1,814	1,674	1,534	1,395	1,256	1,116	977	837	1,541	1,941	1,758	1,610	1,465	1,323	1,186	1,052	921	793	670
San Jacinto	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
San Patricio	0	28	37	46	46	40	34	28	22	17	11	0	0	0	0	0	0	0	0	0	0	0

County	Water Use (AF)										Water Consumption (AF)											
	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060
San Saba	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Schleicher	90	312	468	568	584	507	430	354	277	200	140	61	166	164	195	197	168	140	113	87	61	42
Scurry	10	0	249	341	432	524	466	408	349	291	233	7	0	87	117	146	174	151	130	109	89	70
Shackelford	2	0	156	311	467	421	374	327	280	234	187	2	0	117	226	327	284	243	204	168	134	103
Shelby	1,419	1,658	3,073	2,929	2,785	2,621	2,377	2,133	1,889	1,645	1,400	1,348	1,539	2,771	2,607	2,446	2,270	2,027	1,790	1,561	1,337	1,120
Sherman	2	0	0	92	184	158	132	105	79	53	26	1	0	0	36	69	57	46	36	26	16	8
Smith	5	18	49	80	111	133	118	103	88	74	59	5	17	47	75	101	118	103	88	74	60	47
Somervell	287	184	260	336	413	372	330	289	248	207	165	264	155	195	244	289	251	215	181	149	119	91
Starr	36	48	64	79	79	69	59	49	39	29	18	29	36	40	49	48	41	35	28	22	16	10
Stephens	0	52	184	315	447	402	357	312	268	223	179	0	44	138	229	313	271	232	195	161	128	98
Sterling	57	265	707	881	893	905	765	625	484	344	236	39	141	248	303	302	300	249	199	151	105	71
Stonewall	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Sutton	34	0	390	534	677	821	730	639	547	456	365	23	0	137	183	229	272	237	204	171	140	109
Swisher	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Tarrant	8,805	6,836	5,469	4,101	2,734	1,367	0	0	0	0	0	8,101	5,773	4,102	2,974	1,914	923	0	0	0	0	0
Taylor	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Terrell	10	0	162	221	281	341	303	265	227	189	151	7	0	57	76	95	113	98	84	71	58	45
Terry	3	0	243	332	422	511	454	397	341	284	227	2	0	85	114	142	169	148	127	106	87	68
Throckmorton	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Titus	3	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0
Tom Green	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Travis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Trinity	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Tyler	76	110	147	184	185	161	137	114	90	66	42	72	83	93	114	113	96	80	65	50	36	23
Upshur	4	57	247	437	627	764	690	617	543	469	396	4	54	226	393	555	665	591	519	449	382	316
Upton	1,761	2,955	3,728	3,442	3,156	2,870	2,398	1,927	1,455	983	664	1,198	1,576	1,305	1,171	1,041	916	749	588	433	283	185
Uvalde	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Val Verde	0	0	80	110	139	168	150	131	112	94	75	0	0	28	38	47	56	49	42	35	29	22
Van Zandt	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Victoria	0	35	46	58	58	51	43	36	28	21	14	0	26	29	36	35	30	25	20	16	11	7
Walker	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Waller	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Ward	588	568	683	888	871	855	764	672	581	489	398	114	568	239	297	278	260	228	197	167	138	110
Washington	36	0	497	878	798	718	638	559	479	399	319	29	0	298	516	459	404	351	300	251	204	160
Webb	4,586	3,661	3,476	3,052	2,626	2,244	1,872	1,501	1,128	699	255	3,677	2,627	2,109	1,814	1,529	1,274	1,033	803	580	344	113
Wharton	0	43	57	71	72	62	53	44	35	26	17	0	32	36	44	43	37	31	25	20	14	9
Wheeler	3,792	3,524	3,072	2,620	2,168	1,717	1,265	813	362	0	0	1,896	1,605	1,229	1,015	813	622	443	274	117	0	0
Wichita	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Wilbarger	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Willacy	0	23	31	39	39	34	29	24	19	14	9	0	18	20	24	24	20	17	14	11	8	5
Williamson	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Wilson	417	1,612	1,865	1,679	1,492	1,306	1,119	932	746	560	373	334	1,146	1,119	986	858	734	615	501	392	287	187
Winkler	62	464	618	821	1,024	979	873	767	661	556	450	12	247	216	275	332	305	267	231	195	160	127
Wise	2,314	2,757	2,450	2,144	1,838	1,531	1,225	919	613	306	0	2,129	2,328	1,838	1,555	1,287	1,034	796	574	368	176	0
Wood	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Yoakum	18	238	330	423	384	346	308	269	230	192	154	13	127	116	145	130	115	100	86	72	59	46
Young	8	0	78	157	235	211	188	164	141	118	94	7	0	59	113	164	143	122	103	85	68	52
Zapata	32	41	55	68	68	60	51	42	33	25	16	26	31	35	42	42	35	30	24	19	13	8
Zavala	407	2,065	2,427	2,280	2,167	2,035	1,904	1,773	1,502	1,197	891	326	1,477	1,465	1,351	1,247	1,132	1,020	912	747	575	410
SUM (KAF)	81.5	110	132	135	134	122	104	87	70	53	39	64.8	78.2	76.9	76.0	72.8	64.2	53.2	43.4	34.4	26.3	19.1

Table 16. County-level estimate of 2012-2060 projections for oil and gas water use and water consumption (AF).

County	Water Use (AF)										Water Consumption (AF)											
	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060
Anderson	39	129	140	157	177	181	185	169	147	126	105	67	87	88	98	109	115	121	109	93	78	63
Andrews	3,212	3,481	3,959	3,833	3,710	3,511	3,177	2,842	2,509	2,192	1,929	1,868	1,231	1,029	921	819	742	640	544	453	372	311
Angelina	0	116	220	316	412	427	389	351	312	274	237	32	112	203	286	366	374	336	299	263	228	195
Aransas	0	12	10	8	7	5	5	5	5	5	5	10	11	10	8	6	5	5	5	5	5	5
Archer	30	351	405	444	483	389	344	311	279	246	213	239	326	337	343	344	252	222	194	167	142	119
Armstrong	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Atascosa	1,012	2,993	2,770	2,713	2,706	2,700	2,693	2,393	2,021	1,649	1,279	867	2,155	1,711	1,618	1,551	1,484	1,415	1,219	1,000	790	590
Austin	0	28	127	224	320	288	257	226	194	163	132	20	29	88	143	195	173	151	130	110	91	73
Bailey	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bandera	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bastrop	0	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12
Baylor	1	14	14	14	14	14	13	13	13	13	13	12	12	12	12	12	12	12	12	12	13	13
Bee	66	111	127	129	112	95	80	67	53	40	26	92	90	89	87	74	62	52	44	36	28	20
Bell	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bexar	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Blanco	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Borden	27	272	679	926	927	929	784	639	494	352	244	72	165	263	339	331	323	267	214	164	117	82
Bosque	0	470	557	627	696	579	516	466	416	365	315	238	439	462	485	502	387	340	296	255	216	180
Bowie	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Brazoria	0	91	102	113	125	114	100	88	76	64	52	42	80	78	82	86	76	67	59	52	45	38
Brazos	238	364	741	975	1,207	1,072	938	804	670	536	402	266	286	477	602	721	628	538	451	368	287	211
Brewster	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Briscoe	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Brooks	27	70	77	84	80	69	60	52	44	36	28	50	60	58	60	55	47	41	36	31	26	22
Brown	23	35	34	34	33	32	31	30	29	28	27	17	16	14	14	14	14	14	14	14	14	14
Burleson	247	380	995	1,459	1,923	1,717	1,512	1,306	1,100	892	686	279	299	632	890	1,135	993	855	723	595	471	354
Burnet	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Caldwell	75	98	123	111	98	85	72	59	46	33	20	82	75	81	71	62	54	45	37	29	22	14
Callahan	18	48	52	57	55	47	41	35	30	25	19	34	41	39	41	38	32	28	24	21	18	15
Callahan	84	93	88	88	87	83	79	74	70	66	62	29	24	18	17	16	15	15	15	15	15	15
Cameron	27	58	65	72	68	57	47	39	31	23	15	38	47	45	48	43	34	28	23	18	13	9
Camp	13	12	12	11	11	11	10	9	9	8	8	2	2	1	1	1	0	0	0	0	0	0
Carson	2	14	14	14	14	14	14	14	14	14	14	13	13	13	13	13	13	13	13	13	13	13
Cass	1	30	39	48	58	68	60	52	45	37	30	26	28	36	44	52	60	52	45	38	31	24
Castro	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Chambers	0	9	9	9	9	9	9	9	9	9	9	8	9	9	9	9	9	9	9	9	9	9
Cherokee	10	129	172	216	263	299	267	236	204	173	141	80	123	163	201	239	269	236	204	173	144	116
Childress	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Clay	63	508	613	699	786	655	584	527	471	414	357	318	472	506	538	563	435	382	333	286	243	202
Cochran	56	128	154	181	208	234	210	187	163	139	115	46	64	54	63	71	79	70	62	54	46	38
Coke	520	511	484	480	477	451	425	397	370	346	322	114	84	46	40	33	32	31	29	28	27	26
Coleman	100	113	108	107	107	102	97	91	86	82	77	37	31	24	23	22	21	21	21	21	21	20
Collin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Collingsworth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Colorado	28	129	608	1,078	1,534	1,376	1,221	1,067	913	759	605	86	120	402	667	913	802	697	596	499	406	317
Comal	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Comanche	2	388	444	485	525	419	363	319	276	232	188	224	366	374	379	380	278	240	204	170	138	109
Concho	515	507	480	477	474	448	422	394	367	343	320	114	84	46	40	34	33	31	30	29	28	27
Cooke	1,493	1,708	1,343	978	612	246	28	27	26	25	24	1,391	1,434	1,001	702	421	158	13	13	13	13	13
Conyell	0	569	1,238	1,102	767	434	158	0	0	0	0	236	522	972	827	548	284	103	0	0	0	0

County	Water Use (AF)										Water Consumption (AF)											
	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060
Cottle	32	43	41	41	41	39	38	36	34	33	31	18	16	14	14	13	13	13	13	13	13	13
Crane	280	508	617	728	840	947	861	776	692	610	531	227	246	225	249	273	299	265	232	201	174	149
Crockett	507	1,097	1,732	2,035	1,843	1,552	1,261	971	682	394	207	553	606	641	730	650	539	434	332	235	143	85
Crosby	1,083	1,050	994	987	980	926	871	814	757	706	656	224	161	82	69	55	53	50	47	45	43	40
Culberson	279	293	506	873	1,240	1,535	1,393	1,250	1,110	972	843	151	97	249	308	371	415	368	323	279	240	208
Dallam	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Dallas	79	726	1,076	888	700	512	339	170	0	0	0	134	624	818	651	493	343	220	106	0	0	0
Dawson	268	695	954	1,137	1,164	1,184	1,023	862	703	546	423	165	323	328	371	360	353	296	241	189	140	104
Deaf Smith	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Delta	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Denton	3,249	3,297	2,220	1,136	51	19	13	13	13	13	13	3,108	2,805	1,688	840	44	13	13	13	13	13	13
DeWitt	2,177	2,061	1,858	1,646	1,421	1,188	958	729	500	271	42	1,801	1,493	1,149	999	846	694	550	413	281	155	35
Dickens	0	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	13
Dimmit	3,708	4,874	4,919	5,001	5,001	4,952	4,337	3,580	2,824	2,068	1,315	3,068	3,506	2,980	2,913	2,795	2,648	2,246	1,797	1,368	958	569
Donley	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Duval	52	133	147	160	153	131	114	99	84	69	54	96	114	110	114	105	89	77	68	58	50	41
Eastland	333	578	937	1,091	934	764	644	535	425	318	211	286	303	539	619	472	335	263	196	133	75	21
Ector	845	1,144	1,537	1,612	1,690	1,628	1,435	1,245	1,056	870	725	850	612	588	577	570	520	447	377	310	251	206
Edwards	0	28	29	29	29	29	29	29	29	29	29	28	28	28	29	29	29	29	29	29	29	29
Ellis	38	112	147	180	213	186	164	144	123	103	82	56	99	114	133	150	125	107	90	74	59	45
El Paso	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Erath	12	470	505	521	536	426	376	340	304	268	232	274	443	427	411	391	283	250	218	188	161	134
Falls	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Fannin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Fayette	132	1,149	2,403	2,164	1,887	1,585	1,282	979	677	375	72	166	844	1,476	1,306	1,118	922	733	551	377	210	51
Fisher	432	426	403	401	398	376	355	332	309	289	269	97	71	40	35	30	28	27	26	25	24	24
Floyd	148	156	148	147	146	139	131	123	116	109	102	42	34	23	21	19	19	19	18	18	18	17
Foard	3	12	12	12	12	12	12	12	12	12	11	10	10	10	10	10	10	10	10	10	10	10
Fort Bend	25	66	72	79	75	65	56	49	41	34	27	47	56	54	56	52	44	38	33	29	24	20
Franklin	5	5	5	5	5	5	4	4	4	3	3	1	1	0	0	0	0	0	0	0	0	0
Freestone	429	929	1,117	1,331	1,494	1,458	1,291	1,121	954	785	618	600	844	975	1,133	1,254	1,213	1,056	903	757	615	479
Frio	729	1,167	1,217	1,243	1,250	1,215	1,178	1,142	986	804	620	666	858	772	774	752	702	652	603	504	398	296
Gaines	124	914	1,429	1,846	2,000	1,945	1,671	1,398	1,127	859	651	190	517	590	686	694	635	533	436	344	259	197
Galveston	0	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6
Garza	53	321	395	469	544	491	438	386	334	284	234	44	166	144	164	184	162	142	122	104	87	71
Gillespie	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Glasscock	1,838	2,402	3,057	2,887	2,718	2,355	1,994	1,634	1,275	921	681	1,590	1,165	1,010	923	839	704	575	452	334	224	153
Goliad	25	64	70	77	73	63	54	47	40	33	26	46	55	53	55	50	42	37	32	28	24	20
Gonzales	2,164	1,791	1,600	1,405	1,207	1,010	813	616	418	221	24	1,764	1,288	980	844	712	585	463	346	233	126	23
Gray	68	78	75	74	74	70	67	63	60	57	53	26	22	17	16	15	15	15	15	15	15	15
Grayson	6	19	18	18	18	18	18	18	17	17	17	14	14	14	14	14	14	14	14	14	14	14
Gregg	25	191	274	353	433	476	429	383	337	292	246	71	182	256	322	387	418	371	326	282	240	199
Grimes	95	159	323	483	602	537	471	405	340	275	209	120	129	214	305	367	321	276	233	192	153	115
Guadalupe	0	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
Hale	1,289	1,235	1,168	1,160	1,152	1,087	1,022	954	886	826	766	252	177	82	67	51	48	45	42	39	36	33
Hall	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hamilton	0	361	393	314	236	157	101	51	0	0	0	93	321	312	239	169	103	66	32	0	0	0
Hansford	13	88	577	1,068	904	749	602	456	309	162	16	68	79	261	432	348	278	218	161	108	59	13
Hardeman	0	9	9	9	9	9	10	10	10	10	10	9	9	9	9	9	10	10	10	10	10	10
Hardin	0	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12
Harris	0	24	24	24	24	24	24	24	24	24	24	24	24	24	24	24	24	24	24	24	24	24
Harrison	868	1,763	2,388	2,145	1,956	1,778	1,608	1,438	1,268	1,098	930	1,021	1,658	2,189	1,935	1,735	1,557	1,386	1,219	1,059	903	753

County	Water Use (AF)										Water Consumption (AF)											
	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060
Hartley	7	7	7	7	7	6	6	6	5	5	4	1	1	0	0	0	0	0	0	0	0	0
Haskell	90	98	93	93	92	88	83	79	74	70	66	30	25	18	17	16	16	15	15	15	15	15
Hays	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hemphill	1,441	2,567	2,314	2,037	1,763	1,499	1,244	988	732	476	223	1,498	1,209	971	821	683	562	452	349	252	163	80
Henderson	3	120	176	235	296	346	308	272	235	198	161	91	113	166	218	269	310	272	235	199	164	131
Hidalgo	46	119	130	143	136	117	101	88	74	61	48	85	101	98	102	94	79	69	60	52	44	37
Hill	131	1,572	1,343	1,106	869	632	422	218	13	14	14	244	1,349	1,031	617	427	279	141	13	14	14	14
Hockley	6	18	18	18	18	17	17	17	17	16	16	13	13	13	13	13	13	13	13	13	13	13
Hood	645	529	678	820	961	841	743	651	559	467	375	695	465	528	608	679	566	485	409	338	271	209
Hopkins	42	41	38	38	38	36	34	31	29	27	25	8	6	3	2	2	2	1	1	1	1	1
Houston	178	254	322	287	254	220	187	152	119	85	51	195	196	210	185	161	138	116	94	74	54	35
Howard	619	1,611	2,491	2,939	2,747	2,343	1,940	1,538	1,138	742	476	643	870	898	1,028	938	782	633	490	354	226	142
Hudspeth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hunt	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hutchinson	21	51	156	237	204	173	144	115	86	58	30	32	34	85	110	90	75	62	50	39	28	20
Irion	1,677	2,286	3,192	3,643	3,357	2,890	2,423	1,955	1,487	1,026	713	1,070	937	937	1,065	940	778	621	471	327	190	102
Jack	17	501	575	635	693	572	499	438	378	317	256	232	459	470	487	497	381	328	278	231	187	146
Jackson	25	64	70	77	73	63	55	47	40	33	26	46	55	53	55	51	43	37	32	28	24	20
Jasper	87	118	148	133	118	103	88	73	58	43	28	100	92	98	88	77	67	57	47	38	30	21
Jeff Davis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Jefferson	0	13	13	13	13	14	14	14	14	14	14	13	13	13	13	13	14	14	14	14	14	14
Jim Hogg	33	85	93	102	97	83	72	63	53	44	34	61	73	70	73	67	56	49	43	37	31	26
Jim Wells	25	65	71	78	74	64	55	48	40	33	26	46	55	53	55	51	43	37	33	28	24	20
Johnson	4,192	4,240	3,630	2,809	2,086	1,365	683	10	10	10	10	4,029	3,611	2,690	2,059	1,471	918	447	10	10	10	10
Jones	117	125	119	118	117	111	106	99	93	88	82	35	29	20	19	17	17	17	17	16	16	16
Karnes	3,882	2,820	2,528	2,229	1,919	1,603	1,288	975	662	349	35	3,155	2,028	1,545	1,336	1,127	923	728	542	363	192	29
Kaufman	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Kendall	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Kenedy	42	106	118	130	123	106	92	80	68	55	43	78	92	89	92	85	72	62	55	47	40	33
Kent	29	39	38	38	38	36	35	33	32	31	29	18	16	14	14	13	13	13	13	13	13	13
Kerr	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Kimble	0	17	17	17	17	17	17	17	17	17	17	17	17	17	17	17	17	17	17	17	17	17
King	8,635	8,287	7,836	7,783	7,730	7,293	6,857	6,402	5,946	5,545	5,144	1,704	1,198	565	461	357	334	311	291	271	253	236
Kinney	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Kleberg	27	70	77	84	80	69	60	52	44	36	28	51	60	58	60	55	47	41	36	31	26	22
Knox	3	15	15	15	15	14	14	14	14	14	14	12	12	12	12	12	12	12	12	12	12	12
Lamar	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Lamb	647	620	586	562	579	546	513	479	445	415	385	127	89	41	34	26	24	22	21	19	18	17
Lampasas	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
La Salle	2,889	4,569	4,617	4,705	4,772	4,830	4,263	3,541	2,819	2,098	1,380	2,408	3,293	2,801	2,731	2,647	2,556	2,183	1,757	1,349	959	590
Lavaca	145	1,003	1,613	1,470	1,313	1,148	985	824	662	501	340	179	742	1,005	898	786	673	567	465	368	274	184
Lee	132	230	421	536	650	577	506	435	363	290	218	151	179	272	333	390	340	292	246	201	158	117
Leon	327	847	1,482	1,983	2,481	2,349	2,077	1,802	1,530	1,256	985	361	629	977	1,301	1,611	1,527	1,325	1,129	941	758	584
Liberty	0	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16
Limestone	271	383	402	424	431	391	347	302	257	212	167	356	350	355	363	361	325	283	242	203	166	129
Lipscomb	387	656	1,098	926	758	597	446	294	142	21	21	434	335	467	375	290	221	161	105	52	13	13
Live Oak	1,002	851	814	751	757	776	798	820	729	610	492	853	627	523	473	455	443	433	422	363	294	230
Llano	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Loving	542	691	792	925	1,058	1,020	934	848	762	681	601	300	256	223	251	279	259	229	202	175	152	131
Lubbock	6,211	5,963	5,663	5,673	5,684	5,419	5,089	4,745	4,401	4,097	3,794	1,228	865	433	365	298	290	268	249	229	212	196
Lynn	981	974	1,166	1,246	1,327	1,365	1,255	1,144	1,033	929	826	226	168	179	192	205	227	200	175	150	128	107
McCulloch	42	40	38	38	38	35	33	31	29	27	25	8	6	3	2	2	2	2	1	1	1	1

County	Water Use (AF)										Water Consumption (AF)											
	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060
McLennan	0	194	234	265	296	235	203	177	152	127	101	119	185	197	206	212	154	132	111	91	73	56
McMullen	1,720	2,653	2,912	3,203	3,448	3,666	3,398	3,010	2,622	2,235	1,850	1,465	1,924	1,775	1,860	1,911	1,941	1,746	1,507	1,276	1,066	848
Madison	204	295	597	785	972	861	754	646	538	430	323	227	231	384	485	581	504	432	362	295	231	169
Marion	5	208	348	483	619	622	561	501	440	379	319	73	196	322	438	552	546	485	425	368	312	258
Martin	2,435	2,906	3,527	3,262	2,998	2,657	2,251	1,845	1,441	1,043	771	2,190	1,435	1,191	1,059	933	796	651	513	380	257	177
Mason	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Matagorda	34	87	96	105	100	86	75	64	55	45	35	63	75	72	75	69	58	51	44	38	32	27
Maverick	174	1,652	1,988	2,364	2,737	3,111	2,933	2,617	2,302	1,986	1,674	1,88	1,196	1,201	1,342	1,474	1,597	1,461	1,269	1,085	910	744
Medina	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Menard	1,185	1,148	1,086	1,079	1,071	1,012	952	889	827	772	717	244	175	88	74	59	56	53	50	48	45	43
Midland	1,719	2,876	3,522	3,272	3,025	2,666	2,227	1,788	1,350	918	612	1,661	1,506	1,256	1,127	1,005	855	695	542	395	257	164
Midland	0	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14
Mills	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Mitchell	27	284	361	435	511	460	409	358	309	259	211	50	162	142	163	184	162	141	122	103	86	70
Montague	3,233	3,776	3,228	2,665	2,102	1,538	1,026	525	25	24	24	3,186	3,216	2,452	1,950	1,474	1,025	663	326	14	14	14
Montgomery	0	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15
Moore	4	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16
Morris	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Motley	130	138	132	131	130	123	117	110	103	97	91	39	31	22	20	19	18	18	18	18	17	17
Nacogdoches	1,073	1,642	2,299	2,141	1,986	1,815	1,643	1,471	1,299	1,128	958	1,220	1,550	2,101	1,930	1,764	1,591	1,420	1,251	1,089	932	779
Navarro	11	25	24	24	24	24	23	23	22	22	22	17	16	15	15	15	15	15	15	15	15	15
Newton	98	138	173	156	138	120	102	84	67	49	31	111	107	115	102	89	77	65	54	43	33	23
Nolan	214	218	207	205	204	193	182	171	160	150	140	54	42	26	24	21	21	20	20	19	19	18
Nueces	25	64	70	77	73	63	55	47	40	33	23	46	55	53	55	51	43	37	32	28	24	20
Ochiltree	286	508	824	1,040	853	674	503	332	161	24	23	329	266	355	418	325	247	180	116	57	13	13
Oldham	15	14	13	13	13	12	12	11	10	9	9	3	2	1	1	1	1	1	1	1	1	1
Orange	0	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
Palo Pinto	120	547	656	752	847	709	625	552	480	408	336	281	446	483	524	557	430	370	314	261	212	165
Panola	958	1,578	2,136	1,919	1,749	1,590	1,438	1,286	1,134	983	832	1,095	1,484	1,959	1,731	1,552	1,392	1,240	1,091	948	808	674
Parker	1,083	1,180	1,464	1,733	2,001	1,748	1,545	1,353	1,162	970	779	1,215	1,035	1,139	1,284	1,414	1,176	1,009	851	702	563	434
Parmer	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Pecos	409	543	690	878	1,068	1,180	1,072	966	861	762	672	274	227	313	331	353	359	320	283	249	220	198
Polk	133	195	247	221	195	170	144	118	92	67	41	148	151	162	143	125	107	90	74	58	43	29
Potter	2	14	14	14	14	14	14	14	14	14	14	13	13	13	13	13	13	13	13	13	13	13
Presidio	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Rains	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Randall	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Reagan	1,350	3,414	4,211	3,802	3,395	2,985	2,457	1,931	1,406	886	529	1,361	1,825	1,501	1,323	1,153	991	796	610	432	265	155
Real	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Red River	4	4	4	4	4	4	3	3	3	3	3	1	1	0	0	0	0	0	0	0	0	0
Reeves	611	1,111	1,520	2,067	2,619	2,761	2,522	2,285	2,052	1,827	1,614	701	632	688	796	908	888	791	700	615	541	477
Refugio	23	60	66	72	69	59	51	44	38	31	24	43	51	49	51	47	40	35	30	26	22	18
Roberts	365	1,711	1,502	1,270	1,041	822	611	400	189	20	20	423	819	647	524	412	316	231	151	76	20	20
Robertson	305	691	813	817	826	746	651	556	461	366	271	431	599	654	657	664	595	512	431	354	279	208
Rockwall	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Runnels	285	287	272	271	269	255	240	225	210	197	184	70	53	32	29	26	25	24	24	23	22	22
Rusk	210	719	1,149	1,569	1,994	1,844	1,668	1,492	1,316	1,141	967	323	637	1,017	1,377	1,730	1,578	1,404	1,234	1,070	912	759
Sabine	147	331	584	809	1,035	946	858	770	682	595	508	196	319	536	728	915	826	739	653	571	491	413
San																						
Augustine	1,584	2,198	2,077	1,928	1,779	1,628	1,479	1,330	1,180	1,032	884	1,642	2,052	1,880	1,722	1,567	1,415	1,268	1,124	983	847	715
San Jacinto	0	8	8	8	8	8	9	9	9	9	9	8	8	8	8	8	8	8	8	8	8	8
San Patricio	20	52	57	63	60	51	44	39	33	27	21	38	45	43	45	41	35	30	26	23	19	16

County	Water Use (AF)										Water Consumption (AF)											
	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060	2011	2015	2020	2025	2030	2035	2040	2045	2050	2055	2060
San Saba	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Schleicher	230	473	621	718	732	647	562	477	392	308	241	144	213	199	226	225	194	165	136	109	84	64
Scurry	3	34	280	368	456	544	483	423	363	304	246	37	33	118	143	169	193	168	145	123	102	83
Shackelford	219	464	562	655	747	628	558	500	442	385	328	264	285	329	373	409	305	263	224	187	153	121
Shelby	1,388	1,861	3,283	3,109	2,938	2,754	2,496	2,238	1,980	1,723	1,467	1,536	1,745	2,976	2,781	2,593	2,400	2,143	1,892	1,650	1,414	1,185
Sherman	9	42	35	121	207	178	151	124	98	71	44	28	33	26	55	84	70	59	48	39	29	21
Smith	20	91	107	125	145	163	147	131	115	100	84	67	71	85	100	117	132	117	102	88	74	61
Somervell	287	237	304	367	431	377	333	292	250	209	168	309	208	236	272	304	253	217	183	151	121	93
Starr	35	90	99	108	103	89	77	67	57	46	36	65	77	75	77	71	60	52	46	39	34	28
Stephens	5,158	5,248	5,064	5,103	5,141	4,775	4,458	4,141	3,825	3,541	3,257	1,226	1,004	663	630	591	476	423	374	328	285	244
Stirling	89	343	780	947	953	958	812	667	522	390	270	107	191	290	338	331	325	270	217	166	120	85
Stonewall	629	615	583	579	575	543	511	478	445	416	387	136	99	53	45	38	36	34	33	31	30	29
Sutton	33	59	446	582	720	858	763	668	573	481	389	81	53	185	225	264	303	264	227	192	160	130
Swisher	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Tarrant	8,805	7,084	5,672	4,245	2,817	1,391	12	12	12	12	12	8,313	6,020	4,294	3,105	1,985	935	12	12	12	12	12
Taylor	71	81	77	77	76	73	69	65	62	58	55	26	22	17	16	15	15	15	15	15	15	14
Terrell	502	540	673	724	776	806	740	672	606	544	483	158	128	145	152	160	173	154	136	120	105	92
Terry	90	119	355	439	525	606	543	479	416	354	293	51	45	121	144	168	192	167	144	122	102	83
Throckmorton	200	204	194	193	191	181	171	161	150	141	132	52	40	25	23	20	20	19	19	19	18	18
Titus	8	8	7	7	7	7	6	6	5	5	5	2	1	1	0	0	0	0	0	0	0	0
Tom Green	53	72	69	69	68	66	63	60	58	55	53	31	28	24	24	23	23	23	23	23	23	23
Travis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Trinity	0	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
Tyler	78	123	160	197	198	174	150	127	103	79	55	91	96	106	127	125	109	93	78	63	49	36
Upshur	39	199	379	551	726	851	771	690	609	529	450	95	164	325	474	620	723	644	566	491	419	349
Upton	1,744	3,075	3,887	3,575	3,265	2,960	2,470	1,984	1,499	1,020	699	1,863	1,694	1,458	1,296	1,144	1,001	817	641	473	318	219
Uvalde	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Val Verde	0	66	144	169	195	221	199	179	158	139	120	67	66	91	97	102	108	98	89	81	74	68
Van Zandt	56	65	62	62	61	59	56	53	50	47	45	22	19	15	15	14	14	14	14	14	13	13
Victoria	25	66	72	79	75	65	56	49	41	34	27	47	56	54	56	52	44	38	33	29	24	20
Walker	0	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11
Waller	0	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7
Ward	582	632	775	988	941	915	815	716	617	521	429	622	620	317	362	333	307	267	229	193	161	132
Washington	0	44	545	924	840	757	673	589	506	422	338	30	46	346	561	500	442	385	330	277	227	178
Webb	4,599	3,878	3,708	3,257	2,804	2,397	2,007	1,623	1,238	796	341	3,948	2,844	2,337	2,014	1,701	1,422	1,166	922	687	439	196
Wharton	31	81	89	97	93	80	69	60	51	42	33	58	69	67	69	64	54	47	41	36	30	25
Wheeler	3,794	3,609	3,157	2,682	2,210	1,748	1,293	839	385	22	21	3,850	1,683	1,308	1,071	850	651	469	298	139	20	20
Wichita	59	65	62	62	61	58	55	52	49	46	44	20	17	12	12	11	11	11	11	10	10	10
Wilbarger	7	20	20	20	20	20	19	19	19	18	18	15	14	14	14	14	14	14	14	14	14	14
Willacy	17	44	49	53	51	44	38	33	28	23	18	32	38	37	38	35	29	26	22	19	17	14
Williamson	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Wilson	418	1,671	1,929	1,740	1,548	1,357	1,165	973	782	590	399	373	1,206	1,182	1,045	912	783	659	540	426	315	210
Winkler	152	621	787	977	1,169	1,110	991	873	756	642	531	125	318	295	341	387	351	305	261	220	183	149
Wise	2,313	3,014	2,661	2,293	1,924	1,556	1,238	932	625	319	13	2,348	2,584	2,037	1,691	1,360	1,046	809	587	380	189	13
Wood	17	26	25	25	25	24	23	22	21	21	20	13	12	11	10	10	10	10	10	10	10	10
Yoakum	1,052	1,264	1,300	1,382	1,334	1,240	1,147	1,052	957	870	783	246	299	209	222	191	171	151	132	115	99	84
Young	15	142	197	244	291	236	206	183	159	135	111	125	136	165	188	208	156	135	116	97	81	65
Zapata	30	78	85	93	89	76	66	57	49	40	31	56	66	64	66	61	51	45	39	34	29	24
Zavala	407	2,140	2,531	2,379	2,257	2,118	1,977	1,838	1,559	1,245	932	409	1,555	1,570	1,448	1,336	1,212	1,092	975	802	622	450
SUM (KAF)	118.4	159.3	178.4	179.6	175.1	159.9	139.0	119.1	99.6	81.4	65.4	92.7	96.4	91.8	88.0	71.3	59.7	49.4	39.8	31.3	23.8	18.8




Figure 48. State-level projections to 2060 of HF water use and fresh-water consumption and comparison to earlier water projections.




Figure 49. State-level projections to 2060 of oil and gas industry water use and fresh-water consumption.






Figure 50. Counties with non-zero projected water use. Same coverage as in the 2011 report (thick blue lines) with the addition of Polk, Tyler, Jasper, and Newton counties in East Texas (red circle).




Source: Montgomery et al. (2005)

Source: McMahon and Vaden (2011)

Figure 51. Spatial location of the oil and gas windows in the (a) Barnett Shale and (b) Eagle Ford Shale.




(a)




(b)

Figure 52. Barnett Shale water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.




(a)




(b)

Figure 53. Eagle Ford Shale water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.




(a)




TrackChanges 1.xls

(b)

Figure 54. Pearsall Shale water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.




(a)




TrackChanges 1.xls

(b)

Figure 55. Haynesville and Bossier Shales water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.




(a)




TrackChanges 1.xls

(b)

Figure 56. Haynesville-West Shale water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.




(a)




(b)

Figure 57. East Texas (not including Haynesville and Bossier Shales) water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.




(a)




(b)

Figure 58. Gulf Coast (not including shales) water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.




(a)




TrackChanges 1.xls

(b)

Figure 59. Anadarko Basin water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.



(a)



TrackChanges 1.xls

(b)

Figure 60. Permian Basin water use and consumption projections: (a) comparison with earlier projections; (b) water use and consumption projections under the three scenarios.







Figure 61. Permian Basin water use and consumption projections under the three scenarios: (a) Barnett and Woodford Shales; (b) Wolfcamp Shale and Wolfberry play; and (c) other Permian Basin formations.

V. Conclusions

This update to the 2011 report (whose conclusions were partly summarized in Nicot and Scanlon, 2012) does not fundamentally change the water use projections put forward originally. Both documents outline a water use that is likely to stay in the vicinity of 100 ± 50 kAF/yr for many years. The new projections lower and broaden the expected peak water use and displace the center of gravity of HF water use toward West Texas, an area of the state that has less fresh water. This mechanically translates into a higher brackish water use which when allied with improvement in reuse technologies results in a much lower fresh water consumption than was projected in the 2011 report. The eventual solution in West Texas, after the initial step of using slightly brackish groundwater, is to use more saline brackish water or the abundant produced water from conventional wells to avoid competition with other users who will also rely more and more on brackish water as their water needs increase. In addition to this expected recycling from other uses, the industry itself is making rapidly maturing technological advances that will improve reuse. Fortunately flow back is abundant in most places where fresh water is not (such as in West Texas). However, as in all predictive work, unexpected events can generate large deviations from the projections (as the shale gas revolution did for domestic oil production). The simple discovery of an additional major play (deeper play?) beyond those described in this document could change the state-level water projections. They, however, are unlikely to deviate much in order of magnitude from those outlined here.


It follows that oil and gas water use projections remain a reasonable fraction of mining water use projections, no more than 54% (Figure 62) and a smaller fraction still of the total amount on water use in Texas every year: <0.1 million AF (81.5 kAF in 2011) compared to 15+ million AF (Figure 63).



MiningWaterUse2010-2060_4_TWDB_just.xls

Note: modified from the 2011 report (Nicot et al., 2011, Fig. 135)

Figure 62. Summary of projected water use by mining industry in Texas (2012-2060).



BarPlots_WaterUse_6.xls

Source: TWDB historical water use surveys,

<http://www.twdb.state.tx.us/waterplanning/waterusesurvey/estimates/>

Note: value displayed for mining water use is the 230 kAF from Nicot et al. (2011) rather than the projected 296 kAF listed in TWDB (2012, p.137) or the 2001-2010 average of 184.4 kAF computed with limited information.

Figure 63. Average state level water use (all categories) in 2001-2010.

VI. References

- Bené, P. G., Harden, Bob, Griffin, S. W., and Nicot, J. -P., 2007, Northern Trinity/Woodbine aquifer groundwater availability model: assessment of groundwater use in the northern Trinity aquifer due to urban growth and Barnett Shale development: Texas Water Development Board, TWDB Contract Number 0604830613, 50 p. + apps..
- Fan, L., R. Martin, J. Thompson, K. Atwood, J. Robinson, and G. Lindsay, 2011, An Integrated Approach for Understanding Oil and Gas Reserves Potential in Eagle Ford Shale Formation: SPE 148751.
- McMahon, C., and Vaden, H., 2011, Eagle Ford Shale liquids volumes exceed early expectations. Powell Shale Digest, October 10, 2011, v.1, p. 26-29.
- Montgomery, S. L., Jarvie, D. M., Bowker, K. A., and Pollastro, R. M., 2005, Mississippian Barnett Shale, Fort Worth Basin, north-central Texas: gas-shale play with multi-trillion cubic foot potential. AAPG Bulletin, v. 89, no. 2, p. 155-175.
- Nicot, J. -P., and Potter, E., 2007, Historical and 2006–2025 estimation of ground water use for gas production in the Barnett Shale, North Texas: The University of Texas at Austin, Bureau of Economic Geology, letter report prepared for R. W. Harden & Associates and Texas Water Development Board, 66 p.
- Nicot, J.-P., and Scanlon, B. R., 2012, Water use for shale-gas production in Texas, U.S.: Environmental Science and Technology, v. 46, p. 3580–3586.
- Nicot, J. -P., Hebel, A. K., Ritter, S. M., Walden, S., Baier, R., Galusky, P., Beach, J. A., Kyle, R., Symank, L., and Breton, C., 2011, Current and projected water use in the Texas mining and oil and gas industry: The University of Texas at Austin, Bureau of Economic Geology, Contract Report No. 090480939 prepared for Texas Water Development Board, 357 p. Accessed on 2012: https://www.twdb.state.tx.us/rwpg/rpgm_rpts/0904830939_MiningWaterUse.pdf
- Sinha, S., and Ramakrishnan, H., 2011, A novel screening method for selection of horizontal refracturing candidates in shale gas reservoirs: Society of Petroleum Engineers Paper #144032.
- Texas Water Development Board, 2012, Water for Texas, Vol. II, TWDB Document GP-9-1, January, 392 p.

Appendix 1: Revision to 2011 Report

Although the material below is now obsolete (Table 17), we thought it was important to correct Table 52 of the 2011 report (“Projected water use in the Barnett Shale (Fort Worth Basin”). Although correct values were used in tables of higher order (state level or cumulative across water uses) in the 2011 report, its table 52 was not updated between the draft version and the final version.

Table 17. Update to Table 52 of 2011 report (now obsolete and superseded by this report)

County	2010*	2020	2030	2040	2050	2060
	AF					
Archer	0	1,618	1,292	369	0	0
Bosque	913	2,547	1,065	0	0	0
Clay	634 951	3,734 5,596	1,663 2,495	0	0	0
Comanche	429	2,524	1,125	0	0	0
Cooke	101	282	118	0	0	0
Coryell	0	1,793	1,140	263	0	0
Dallas	620	769	271	0	0	0
Denton	1,674	587	0	0	0	0
Eastland	0	1,127	1,157	386	0	0
Ellis	325	235	63	0	0	0
Erath	2,017	2,500	882	0	0	0
Hamilton	190	1,118	498	0	0	0
Hill	1,008	1,249	441	0	0	0
Hood	1,720	990	215	0	0	0
Jack	1,835 2,386	1,706 2,218	535 696	0	0	0
Johnson	3,308	1,537	241	0	0	0
McLennan	0	1,380	680	62	0	0
Montague	539 809	3,174 4,760	1,415 2,122	0	0	0
Palo Pinto	446	2,627	1,171	0	0	0
Parker	4,003	1,787	153	0	0	0
Shackelford	0	1,121	1,151	384	0	0
Somervell	771	443	96	0	0	0
Stephens	0	1,854	1,178	272	0	0
Tarrant	3,147	1,104	0	0	0	0
Wise	4,220 4,642	1,964 2,157	308 338	0	0	0
Young	0	563	578	193	0	0
Total (Th. AF)	27.9 29.5	40.3 44.5	17.4 19.2	1.9	0.0	0.0

Note: double strikethrough on the incorrect values replaced by the correct but obsolete values.

